Plant Disease Reporter

Agricultural Research Center

Agricultural Research Service

UNITED STATES DEPARTMENT OF AGRICULTURE

VOLUME 57 NUMBER 13

PLDRA4 57(13) 1073-1110

1973 Index
(Issued May 1974)

The Plant Disease Reporter is published monthly as a service to American agriculture. The articles emphasize new developments in plant pathology, including disease incidence and severity, serious outbreaks and epidemics, investigative techniques, and control chemicals. The rapid issuance of such information permits plant pathologists to recommend the judicious use of pesticides or other appropriate control measures against impending disease epidemics. Agricultural Research Center, in publishing this material, does not assume responsibility for the subject matter.

PLANT DISEASE REPORTER INDEX

Issued by

APPLIED PLANT PATHOLOGY LABORATORY
Plant Protection Institute

Beltsville Agricultural Research Center
Volume 57

Beltsville, Maryland
Number 13

AUTHOR INDEX

AAPOLA, A. A., 552
AGARWALA, R. K., (361)
AGBOOLA, S. D., (824)
AHMAD, MUSHTAQ, 800
AHMAD, RIAZ, 505
AHMED, JASIM M., 855
ALASOADURA, S. O., (375)
ALCONERO, R., (884)
ANAGNOSTAKIS, S. L., (158), 225
ANDERSON, N. A., (890)
ANDERSON, ROBERT L., 254
ANDERSON, W. C., (164)
ARTMAN, J. D., 182
AUSHER, R., (53)
AYERS, J. E., (253), (411)
AZERI, TURHAN, 149

BAIN, D. C., (507)
BARKER, K. R., (750), 1025
BARKSDALE, T. H., 964
BARNETT, O. W., 917
BARNETT, PAUL E., 672
BATSON, WILLIAM E., Jr., 453
BAXTER, LUTHER W., (315), (917)
BECKMAN, C. H., 928
BELL, D. K., 90, 549
BeMILLER, J. N., (308), (1043)
BERAHA, L., 445
BERG, C. C., (563)
BERGER, R. D., 837
BERGGREN, G. T., 642
BERGQUIST, R. R., 272
BEUTE, M. K., (387), (1035)
BHALLA, R. B., 6
BIEHN, W. L., 35, 37
BIELENBERG, O., (353)
BIRCHFIELD, W., (199)
BIRD, G. W., 399, 402
BISSONNETTE, H. L., (353)
BLAHA, G., (529)
BLANCO, M., (253), (411), (822)
BLAZQUEZ, C. H., (921)
BLOOM, J. R., (891)
BLOUNT, V., (724)
BOCKHOLT, A. J., (42)
BOESEWINKEL, H. J., 136
BOOSALIS, MICHAEL G., (681), (873)
BORECKA, H., 31
BORECKI, Z., (31)
BRASHEARS, A. D., (129)
BRASHER, E. P., 542
BRAUN, ALVIN J., 431, (433)
BRAVERMAN, S. W., (592), (709)
BRENNAN, EILEEN, (1023)
BRIDGE, JOHN, 798
BRIDGMON, G. H., (576)
BRODRICK, H. T., (173)
BROWN, J. F., (701), 856
BRUEHL, G. W., (935)
BUCHENAUER, H., 460
BUGBEE, W. M., 204
BURGIS, D. S., (78), (119), (724)
BURNS, E. E., 630
BURTON, CLYDE, (62)
BURTON, GLENN W., (262)
BUSBICE, T. H., (946)
BYTHER, R. S., 966

CALPOUZOS, L., (99)
CALVERT, OSCAR, H., (254), (937)
CAMERON, H. R., 241
CAMPBELL, J. M., (932)
CAMPBELL, W. P., 735
CAPOOR, S. P., 160
CARNES, D., (529)
CARVER, R. B., (16)
CEPONIS, M. J., 487
CHALFANT, R. B., (684)
CHAMBERS, ALBERT Y., (848)
CHAMBLEE, D. S., (946)
CHANEY, D. H., 192
CHIDAMBARAM, PALANI, 935
CHIKO, ARTHUR W., 639
CLAFLIN, L. E., 52, 701
CLARK, C. A., 210
CLAYTON, J. L., (187)
COHN, E., 53
COLE, H., Jr., 891
COOK, G. E., 873, (1070)
COOLEY, THOMAS N., (234)
CORDAS, D. I., 926
COX, E. A., (177), (179)
COYNE, DERMOT P., (74), 111, (1070)
CRAWFORD, JOHNNY L., (399), 403
CRILL, PAT, 78, 119, (122), 724, 921
CROSBY, F. L., 244, (329)
CUNNINGHAM, JOHN L., 793

DAFT, GILBERT C., 156
DALE, J. L., 711
DALMACIO, S., (253), (411), (822)
DANIEL, J. T., (695)

DAVIDSON, ROY M., Jr., (522)
DAVIS, J. R., 803
DAVIS, ROBERT E., 333, (987)
DAVIS, SPENCER H., Jr., 28
DAVISON, ARLEN D., 522
DEMSKI, JAMES W., 978
DERRICK, K. S., (16)
DHINGRA, O. D., 185
DOGGETT, C. A., (325)
DOLAN, D. D., (468)
DOOLING, OSCAR J., 298, (701)
DOVAS, C., (321)
DOW, ALICE T., (896)
DRIVER, CHAS. H., (167)
DUBOS, BERNADETTE, (25)
DUNKLE, LARRY D., (681), (873)
DUNLEAVY, J. M., 660
DUNN, R. A., 728
DUTT, KESHAV, (361)

EDMONDS, ROBERT L., 1058
EDMUNDS, J. E., (678)
EDWARDS, D. I., 47, (207)
ELDRIDGE, B. JOE, (169)
ELGIN, J. H., Jr., 657
EL-GOORANI, M. A., 646
ELLETT, C. WAYNE, (555), (807)
ELLINGBOE, A. H., (187)
EMMATTY, D. A., 584, 787
ENGELHARD, A. W., (181)
EPSTEIN, A. H., (471)
ERDMANN, GAYNE G., (230)
ERWIN, D. C., (190), (460)
ESSER, R. P., 558, 1045
EVANS, D. W., (657)

FAJOLA, A. O., 375
FARLEY, J. D., 767
FARRELL, K. M., (260)
FASSULIOTIS, GEORGE, 606
FAULKNER, L. R., (657)
FEDER, W. A., (774)
FENSTERMACHER, J. M., (84)
FICK, G. N., (524)
FIELDHOUSE, D. J., (542)
FILER, T. H., Jr., (59), 338, 341
FILIP, GREGORY M., 499
FILSINGER, ELIZABETH C., (664), (756), (828)
FISHER, J. W., (660)
FLOWERS, R. A., 190, 832
FORER, L. B., 104, (891)
FOX, W. T., (695)
FRAME, ROBERT E., (1055)
FREDERIKSEN, R. A., 42
FREEMAN, T. E., 601
FRENCH, W. J., 902
FRIDLUND, P. R., 483, 865
FRY, GAYLE, (478)
FULTON, J. P., 153, (1029)

GABRIELSON, R. L., 164
GARDNER, WAYNE S., 106
GARNSEY, S. M., 886, 1010
GARRAWAY, M. O., 518
GAY, J. DAN, 13, 684
GEORGOPOULOS, S. G., 321
GESSEL, T. G., (942)
GIBBS, A. F., 544
GILL, C. C., 862
GILL, D. L., 739
GILLASPIE, A. G., Jr., 987
GILPATRICK, J. D., 457, (817)
GOLDEN, A. M., (946)
GOLLIFER, D. E., 701
GOOD, J. M., (169)
GOOD, R. L., 981
GOODING, G. V., Jr., 200, (551), (841), (845)
GOTAN, S. M., (192)
GOWEN, S. R., 678
GRAND, L. F., 325
GRAY, F. A., (657)
GRAY, L. E., 853
GREENBERGER, A., (125)
GRENDAHL, JAN, (814)
GRIFFIN, G. D., 942
GRIFFITHS, D. A., (735)
GRIMM, G. R., 669
GROSCLAUDE, C., 25
GUPTA, G. K., 361

HAGEDORN, D. J., (248), (349)
HALFON-MEIRI, ALIZA, 960
HALL, D. H., (1066)
HALLIWELL, ROBERT S., (81)
HAQUE, S. Q., 412
HARDER, PAUL R., 924
HARE, ROBERT C., 442, 776, 1013
HARIDASAN, M., (1050)
HARMAN, G. E., 592, 709
HARPAZ, I., (125)
HARRINGTON, J. B., Jr., 937
HARRIS, C. M., 44
HARRIS, HASKELL, (549)
HARTILL, W. F. T., 932
HARVEY, J. M., (44)
HARWOOD, ROBERT F., (800)
HAWARE, M. P., 436
HAWKSWORTH, FRANK G., (904)
HEATHER, WILLIAM A., (1058)
HEINZ, D. J., (966)
HEIT, C. E., (592)
HELBURG, LAWRENCE B., (511), 513
HEMSTREET, C., (416)
HENDRIX, JAMES W., (190)
HENNEN, JOE F., (262)
HESKETH, J. D., (1050)
HIBBEN, CRAIG R., 396
HILL, J., (253), (411)
HILL, J. H., 471
HITE, R. E., 131, 760

HOFF, BETTY, (74)
HOFFMANN, J. A., (1006)
HOGGER, Ch., (402)
HOLCOMB, G. E., 16
HOOKER, A. L., 195, 586, (590)
HORN, N. L., 811
HORSFALL, J. G., (276)
HOYLE, M. C., (499)
HSIEH, S. P. Y., (537)
HUDLER, GEORGE W., (55)
HUGHES, J. E., (244), (329)
HUNTER, J. E., (86), (907)
HURLIMANN, J. H., (115)
HUSSEY, R. S., 1025
HUTCHISON, D. J., (669)
HYRE, R. A., 627

IONNAIDIS, N. M., 39
ISHII, M., 475

JACKSON, NOEL, 84
JACOBS, C. J., 173
JACOBSEN, B. J., (890)
JATALA, PARVIZ, 1021
JAWORSKI, C. A., (974)
JAYNES, R. A., (225)
JENSEN, H. J., (1021)
JENSEN, KEITH F., 914
JENSEN, R. L., (811)
JODON, MILDRED H., (891)
JOHN, C. A., (584), (787)
JOHNSON, A. W., (684), (832)
JOHNSON, HENRY E., (948)
JOHNSON, JERRAL D., 81
JOHNSON, LEANDER F., 848
JONES, ALAN L., 62, 428
JONES, JOHN PAUL, (78), (119), 122, 612, (724)
JOSHI, L. K., (531)
JOSHI, L. M., 8
JUDD, R. W., Jr., 624

KAISER, WALTER J., 742
KAMENSKY, SUSAN A., 230
KANG, M. S., (308)
KAPUR, S. P., 598
KAUFFMAN, H. E., 537, (784)
KAUFMAN, J., (487)
KEPLINGER, J. A., (431), 433
KHAN, INAM ULLAH, (505)
KIM, K. S., (153), 1029
KIM, SOON KWON, (586)
KIMBROUGH, J. W., (181)
KINMAN, M. L., (524)
KINSEY, J. G., (344), 590
KISSLER, J. J., 115, (619)
KLEIN, M., 125
KLIEJUNAS, J. T., 383
KNAUSS, J. F., 222, (504)
KNUTSON, DONALD M., (716)
KO, W. H., (383), 907
KOMMEDAHL, THOR, (912)
KOUCHEKI, H. K., 284, 909
KRUPINSKY, J. M., (363)
KUNIMOTO, R. K., (907)
KYROU, N. C., 1032, 1033

LACY, M. L., 425
LADD, S. L., (966)
LAEMMLEN, F. F., (215), (217)
LAI, MINGTAN, 258
LAIRD, PETER P., 94
LAPP, N. A., 551
LARSEN, P. O., 76
LAUT, JOHN G., 511
LEAPHART, CHARLES D., 948
LEBEN, CURT, (156), (515)
LEPPIK, E. E., 704
LEWIS, J. A., 876
LIDER, J. V., (115)
LIM, S. M., (195), 344
LINDBERG, G. D., 420
LINE, ROLAND F., 1006
LOCKE, BOBBY J., (90)
LOEGERING, W. Q., (6)
LONGENECKER, JOHN, (984)
LOONAN, DANIEL V., (780)
LORBEER, J. W., (210)
LUCAS, L. T., 946
LUKENS, R. J., 276
LUKEZIC, F. L., (104)
LUVISI, D. A., (619)

MABBETT, T. H., 294
MAGIE, R. O., (957)
MAHAPATRA, B. K., (278)
MAHMOOD, M., (495)
MAI, W. F., (728)
MAIN, C. E., (39)
MANNING, W. J., 177, 179, 774
MARBUT, SAMUEL A., (315)
MARSH, PAUL B., 664, (756), (828), 994
MARTIN, W. J., 199
MASSIE, L. B., 730
MATUSZEWSKI, MARK, 17
MAY, JAMES, (358), (359)
MAYAMA, M., (308), (1043)
MAYAMA, S., (308), (1043)
McCAIN, A. H., (67)
McCARTNEY, WILLIAM O., 904
McCOY, R. E., 33
McELROY, F. D., 492
McGLOHON, N. E., (399)
McLAUGHLIN, M. R., (471)
McRITCHIE, J. J., 181, 389, (500)
McVEY, D. V., (880)
MERCA, S. D., (537)
MERRILL, WILLIAM, 984
MERWARTH, FRED L., (697)
MESTERHAZY, AKOS, (195)
MICHELL, R. E., 207

MILBRATH, J. A., (241)
MILLER, D. L., 346
MILLER, J. W., 500, 504
MILLER, P. M., 158, 267, 438, 676
MILLER, T. D., (767)
MILLIKAN, D. F., (31)
MINK, G. I., (552)
MINTON, N. A., 946
MOHAMMAD, ABU, 495
MOLINE, H. E., 373
MOLLER, WILLIAM J., (192), (634)
MOORE, B. J., 311
MOORE, B. SHAIN, (822)
MOORE, LAURENCE D., (578)
MORGAN, PAUL, 167
MORROW, RONALD A., (513)
MOSSAHEBI, G. H., (742)
MUIR, J. A., 951
MULDER, N. J., (173)
MULLANY, R., (276)
MULLER, R. A., (529)
MUMFORD, PAULINE, (308)
MURDOCH, C. L., 215, 217
MURPHY, JUDITH A., (308)
MYHRE, D. L., 1050

NARAIN, A., 278
NELSON, PAUL E., (891)
NELSON, R. R., 18, 145, 253, 304, 411, (730), 822, 971, 1047
NENE, Y. L., 463, (572)
NEWSOM, L. D., (811)
NEWTON, MICHAEL, (94)
NICHOLLS, THOMAS H., 55, 263
NICHOLSON, J. F., (185), 269, 531, 770
NICHOLSON, R. L., (143)
NICKEL, JOHN L., (414)
NIELSEN, L. W., 371
NJOMOU, S. E., (529)
NORSE, D., 404, 582
NWIGWE, CHINENYE, 227, 955
NYLAND, GEORGE, 634
NYVALL, R. F., (164), (471)

O'BRIEN, JAMES T., 246
ODVODY, GARY N., 681, (873)
OITTO, W. A., (20)
OKHOVAT, M., (742)
OLIARI, LOURENCO, 870
ORELLANA, R. G., 318
ORR, CALVIN C., 129
OSHIMA, R. J., 719
OSWALD, T. H., 789
OSWALT, T. W., (391)

PALMER, L. T., (8)
PAPPELIS, A. J., 308, 1043
PAPPELIS, G. A., 308, (1043)
PARASRAM, S., (412)
PARTHASARATHY, M. V., 861
PARTRIDGE, A. D., (346)
PARVIN, P. E., (215), (217)
PATEL, B. N., 507
PATIL, S. S., 86
PEADEN, R. N., (657)
PEARSON, R. C., 1066
PEIRSON, D. Q., (603)
PEÑA-IGLESIAS, ANTONIO, (649)
PERKINS, I., (774)
PERSLEY, GABRIELLE J., 1040
PETERSON, J. L., (28)
PFLEGER, F. L., 468, (592), 595, (709)
PHELPS, RALPH H., (294)
PHILLIPS, DOUGLAS J., 814
PHILLIPS, D. V., (371), 1063
PINCKARD, J. A., (642)
PIRONE, T. P., 841, 845
PITKETHLEY, R. N., 803
PITRE, H. N., (533), (1050)
POE, S. L., (921)
POEHLMAN, J. M., (278)
POLLACK, FLORA G., 296
PONCHILLIA, PAUL E., 489
PONTIS, R. E., 379
POWELL, C. C., 515, (603)
POWELL, J. M., 283
PRABHU, A. S., 1000
PRAKASH, VED, (1000)
PRASAD, KRISHNA, 795
PROVVIDENTI, R., 280, 688

RAABE, R. D., (115)
RANNEY, C. D., 325
RASKI, D. J., (115), 416, 619
RATCLIFFE, T. J., (974)
RAVENSCROFT, A. V., (187)
REBOIS, R. V., 169
REDDY, A. P. K., (537), 784
REID, J. T., 329
REXRODE, CHARLES O., 1055
RHOADS, ANN, 1023
RICARD, J., (25)
RICH, A. E., (499)
RICH, S., (158), (438)
RICHARDSON, JEAN, (414)
RICHARDSON, LLOYD T., 3
RIEDEL, R. M., 603
RIFFLE, JERRY W., 355
ROBBINS, R. T., 750
ROBINS, J. K., (951)
ROELFS, A. P., 434, 754, 880
ROTHMAN, P. G., (754)
ROWE, R. C., 387, 1035
ROWELL, J. B., (434), 567, 653
RUBIO-HUERTOS, MIGUEL, 649
RUSH, M. C., 715

San ANTONIO, J. P., 724
SANDS, D. C., (438)
SANTAMOUR, FRANK S., Jr., 997
SASSER, M., (542)
SCHAREN, A. L., 363

SCHARPF, ROBERT F., (904)
SCHENCK, N. C., (981)
SCHLÖSSER, E., (615)
SCHMITT, R. V., (115), (416), (619)
SCHNEIDER, R. W., (824)
SCHOMAKER, MICHAEL E., (511), (513)
SCHUSTER, M. L., 74, (111)
SEARS, E. R., (6)
SECHLER, DALE T., (278)
SEGALL, R. H., 896
SEKHON, H. S., 576
SELA, I., (125)
SELF, R. L., (1072)
SETLIFF, E. C., 473
SHARMA, N. D., (436)
SHARVELLE, E. G., (143)
SHAUNAK, K. K., 533
SHEARER, B. L., 99, 367
SHERWOOD, R. T., 563
SHIGO, A. L., (499)
SHIMABUKURO, R. A., (1021)
SHURTLEFF, M. C., (630)
SIKORA, R. A., 141, 615
SIMPSON, MARION E., (664), 756, 828, (994)
SINCLAIR, J. B., (185), (269), (531), (770), (853)
SINGH, A., 572
SINGH, KISHAN, 220
SINGH, N. D., 260
SINGH, R. S., (141)
SINHA, S. K., (278)
SITARAMAIAH, K., (141)
SKILLING, DARROLL D., (55)
SLEESMAN, JOHN P., (76), 555, (807)
SMILEY, J. H., (841)
SMITH, D. H., (244), (329)
SMITH, D. R., (195)
SMITH, N. A., (187)
SMITH, R. J., Jr., 695
SMITH, RICHARD S., Jr., 67, 69
SMYLY, W. B., 59
SONODA, RONALD M., 260, 638, 747
SRAGO, M. D., (67)
STADELBACHER, GLENN J., (795)
STATLER, G. D., 449
STEADMAN, J. R., 1070
STEINER, G. W., (966)
STENBERG, D., (291)
STEVENS, H., (291)
STEVENSON, JOHN A., (296)
STEVENSON, W. R., 248, 349
STIENSTRA, WARD C., (300), 890
STONE, ERIC G., 884
STONER, A. K., (169), (964)
STRIDER, D. L., 969, 1015, 1020
STROBEL, J. W., (78)
STUTEVILLE, D. L., (52)
SUBIRATS, F. J., 1072
SURIN, PREECHA, (833)
SWARTS, H. D., (173)
SYLVESTER, EDWARD S., 414
SZKOLNIK, MICHAEL, 817

TANG, C. S., (86)
TATE, L. A., (241)
TATTAR, TERRY A., 338, (394)
TAYLOR, G. S., (267)
TEGETHOFF, ALFRED C., 407
TEILLON, H. BRENTON, (338)
TEMPLETON, G. E., (695)
TERRY, E. R., 509
THAPLIYAL, P. N., (269)
THIRUMALACHAR, M. J., (160)
THOMAS, H. L., (544)
THOMPSON, JOHN H., 394
THOMPSON, L. S., (237)
THOMPSON, SAMUEL S., (90)
THYR, B. D., 974
TIETJEN, W. H., (487)
TIMMER, L. W., 899
TINNIN, ROBERT O., 716
TOLER, R. W., (16)
TOLIN, SUE A., (200)
TROIANO, JOHN, (1023)
TROLL, JOSEPH, (924)
TUCKER, E. T., (946)
TUNG, GABRIELLE, (145), (304), (730), (971), (1047)

ULLSTRUP, A. J., (42), (143)
UYEMOTO, J. K., (280)

van der MERWE, J. J. H., 1005
van der ZWET, T., 20
Van SICKLE, G. A., 608, 765
van WYK, P. S., (1005)
VARDARO, P. M., (177)
VIDAVER, ANNE K., (291)
VIEIRA, CLIBAS, (870)

WADE, E. K., (473)
WAGNER, V. R., (104)
WALDHER, J. T., (1006)
WALKER, E. BRADFORD, (338)
WALLIN, JACK R., 298, 780
WALTERS, H. J., 833
WALTON, G. S., (624)
WARREN, H. L., 143, 912
WATKINS, J. E., (449)
WATSON, TOM, (258)
WEATHERS, L. G., (598)
WEBB, RAYMON E., (974)
WEHUNT, E. J., (47)
WELLS, HOMER D., 262, (549)
WELLS, J. C., (387)
WELLS, JOHN M., 234, 697
WELSH, MAURICE F., 358, 359
WHIDDEN, R., (886), (1010)
WHITE, DONALD G., (76), (555), 807
WHITESIDE, J. O., 391, 691
WICKS, T., 560

WIESE, M. V., 187
WILCOXSON, ROY D., 353, (544)
WILKINSON, N. W., (283)
WILKINSON, R. E., (870)
WILLIAMS, R. J., 824
WILLIS, C. B., 237
WILLS, W. H., 578
WITCHER, WESLEY, 315
WOLTZ, S. S., 957
WORLEY, J. F., (987)
WRIGHT, W. R., (445)
WYLLIE, T. D., 478, (789)
WYSONG, D. S., 291

ZEIDERS, K. E., (563)
ZENTMYER, G. A., 529
ZEYEN, R. J., 300
ZIMMER, D. E., 524, 647
ZITTER, T. A., 991

SUBJECT INDEX

Abies balsamea (balsam fir): a quantitative survey for needle rust, 765
--- grandis (grand fir): Poria weirii, Fomes nigrolimitatus, or Armillaria mellea were the most common root-rotting fungi found in dying or discolored fir in Idaho, results indicate root rot is incited by fungus complexes more often than by one organism, 346
Acyrothosiphon pisum: 552; 688
Aerial detection: of tobacco ringspot virus-infected soybean plants in Iowa, 471
Aerial photographs: to detect infestation of jack pine by dwarf mistletoe, 951
Aerial spray: of benomyl provided long term effective control of Sclerotinia sclerotiorum in cabbage seed fields, 164
Africa: 414
Age: effect of age on virulence of Septoria nodorum spores on wheat, 363
Air pollution: tobacco plants in Brookings, South Dakota showed injury from ozone, 106; ozone injury in potato variety trials in Delaware, 542; ozone, reduction in plant size and fresh and dry weights of ears in 'Golden Jubilee' corn, from exposure to ozone, 719; five poinsettia cultivars show considerable resistance to ozone injury, 774; ozone fumigation of 9 forest trees, 914; injury to young buds of Easter lily with ethylene gas in experimental chamber, 1023
Alabama: 296
Alder (Alnus glutinosa): seedlings showed ozone tolerance, 914
Alfalfa: bacterial leaf spot, survival of Xanthomonas alfalfae in alfalfa debris and soil, 52; alfalfa hay yield increased by control of foliar diseases, 353; field resistance to Pseudopeziza medicaginis, 468; 600-800 larvae per plant appeared optimum for screening alfalfa seedlings for resistance to root-knot nematodes, 657; new alfalfa variety Apalachee showed resistance to Stemphylium leafspot, 946
Allium cepa (see Onion)
Alternaria sp.: 756
--- alternata: 39; 592
--- brassicicola: 704
--- crassa: 960
--- solani: 964
--- tenuis: 487
--- tenuissima: 1072
--- triticina: 1000
Anthracnose: of crownvetch, Colletotrichum destructivum isolated from diseased crownvetch in Pennsylvania and was pathogenic, 104; of sweet corn, C. graminicola occurred in epiphytotic proportion in 1972 in Benton County, Indiana, 143; of mango, caused by C. gloeosporioides, controlled with benomyl, 173; of strawberry, caused by C. dematium, 445; of blueberries from New Jersey, 487; a strain of C. gloeosporioides, pathogenic to northern jointvetch, a rice field weed, is endemic in rice areas of Arkansas, 695; physiologic races of C. lindemuthianum, in Minas Gerais, Brazil, 870
Antibiotics (see also Chemicals, Fungicides, and Nematicides): Polyoxin, Kasumin and Pomycin, from species of Streptomyces were ineffective in control of gray mold of strawberries, 31; cycloheximide controlled needlecast on red pine seedlings under nursery conditions, 263
---: streptomycin sulfate: not effective for control of diseases of redcedar and Arizona cypress, 315; ineffective for control of bacterial canker of tomato, 767
---: tetracycline: tetracycline hydrochloride, spraying or injecting lime plants reduced greening symptoms, but disease reappeared within 60 days, 160; tetracycline and oxytetracycline effectively injected into elm trees with a new pressure injector, 338; suppression of elm phloem necrosis symptoms by injection of tetracycline antibiotics, 341; control of pear decline with oxytetracycline HCl by transfusion of 6 to 8 qt of antibiotic per tree, 634; ineffective for control of bacterial canker of tomato, 767

Aphanomyces euteiches: 876
Aphelenchoides cibolensis: 355
--- composticola: 355
--- fragariae: 603; 1015
Aphids: under experimental conditions no evidence of transmission of barley stripe mosaic virus by aphids, leafhoppers or grasshoppers was observed, 639
Aphis fabae: 533
--- gossypii: 533
Apioporthe anomala: 522
Apple: fungicide-supplemented wound dressings for control of Botryosphaeria ribis, 28; leaf pucker disease severity of symptom expression appears to be determined by temperature during the 40-day full bloom and post-bloom period, 358; blossom symptoms associated with three virus diseases of apple, 359; sudden decline and death of apple trees in north-central Wisconsin involved Stereum purpureum, but the silvery leaf symptom was absent, 473; scab, Venturia inaequalis tolerance to dodine in apple orchards in New York, 817; chlorotic leaf spot virus, distribution in budsticks was greatest in the basal end, 865
--- frog eye leaf spot: caused by Physalospora obtusa, covering or fertilizing apple brush piles reduced production of conidia, 158; cultivars differ in susceptibility to P. obtusa, 676
Application time: of fungicide BT for best control of wheat leaf rust, 653
Apricot: apricot leaf curl reported for first time from India is caused by Taphrina deformans, 361
Arborvitae (see Thuja orientalis)
Arceuthobium americanum: 298; 951
--- douglasii: 716
Argentina: 379
Arkansas: 153; 695; 1029
Armillaria mellea: 155; 346; 355
Ascochyta gossypii: 403
--- piniperda (see Sirococcus strobilinus)
--- rabiei: 742
Asexual reproduction: of Botrytis cinerea, the near-ultraviolet transmission properties of assorted glass and plastic materials were closely associated with amounts of sporulation of the fungus beneath the filters, 131
Ash (see Fraxinus)
Aspergillus flavus: 185; 664; 756
--- niger: 756
Assay hosts: of Hydrangea ringspot virus, 300
Australia: 367; 560; 735; 803; 1040; 1058

Bacilliform particles: in cortex cells of Cucumis melo fruits, 649
Bacteria: two bacterial isolates, a pseudomonad, a xanthomonad and a Fusarium isolate found to be involved in a disease of Pelargonium hortorum, formerly attributed to Verticillium albo-atrum, 284
Bacterial diseases: hosts of Xanthomonas albilineans, 1040
---: bacterial blight: of bean: virulent strain of Xanthomonas phaseoli from Uganda compared with strains from Nebraska and Colombia, 74; several lines and cultivars of Phaseolus showed tolerant reaction to Nebraska strains of X. phaseoli, 111
---:---: of Gerbera jamesonii, incited by Pseudomonas cichorii in Florida, 504
---: ---: of soybeans: Pseudomonas glycinea capable of overwintering in Ohio but survival rate is low, 156; found seedborne in 27 seedlots from five districts in India, first report of this bacterium occurring in soybean in India, 531
---: bacterial canker: of tomato: new tomato variety H2990 shows high resistance, 584; disease can be spread during clipping of transplants, 767; was eradicated from infected tomato seed by seed extraction with HCl and drying for 3 hr at 66°C in a tumble drier, 974
---: bacterial leaf blight: of rice: persistence of Xanthomonas oryzae in the infected tissues of rice plants, 227; a leaf clipping technique to evaluate susceptibility of rice varieties to X. oryzae, 537; initial infection in inoculated resistant and susceptible cultivars was equal, but some unknown mechanism in resistant plant caused slower movement and multiplication of the bacterium, 784
---: bacterial leaf spot: of alfalfa, virulent X. alfalfae was recovered from infested alfalfa debris and naturally and artificially infested soil monthly from August 1971 to June 1972, in Kansas, and in infected alfalfa shoots stored for 8 years in a laboratory, 52; apparently the first report of bacterial spot of tomato and pepper in California, caused by X. vesicatoria, 258; of zinnia, caused by X. nigromaculans f. sp. zinniae, in Ohio, 555; found on zinnia in North Carolina, 1020
--- bacterial wilt: of carnation, in New Zealand, 136; of cassava, 824; Goss's bacterial wilt, an undescribed Corynebacterium is cause of a foliar blight on corn in Nebraska, 291
---: blackleg of potato, control with seed and soil treatments, 803
---: fire blight: of pear: a method for screening seedlings for resistance, 20;

Egypt has been free of the disease for the last 6 years, 646
---: leaf scald of sugarcane, a quantitative inoculation technique for screening sugarcane varieties for resistance, 582
---: soft rot of carrots, level of bacterial contamination and postharvest holding temperature affect incidence, 896
Bacterium: a small, thin, rod-shaped, sometimes bent bacterium is consistently associated with ratoon stunting disease of sugarcane, 987
Ball moss (Tillandsia recurvata): control of this epiphytic plant in Texas, with certain metal-containing fungicides, 81
Balsam fir (see Abies)
Banana: decline in Lebanon thought to be associated with nematodes and fungi interaction, 615; extraction method for estimating nematode populations in banana roots, 678
Barbados: 404
Bark beetle: Pseudopityophthorus sp., effect of girdling or felling oak-wilted trees on oak bark beetle attacks in West Virginia, 1055
Barley: 16 of 73 Indian and American barley cultivars showed immunity or high resistance to seven isolates of Helminthosporium gramineum in India, 495; found to be new host of H. maydis and is added to host range of race 2 of H. carbonum, 507; barley stripe mosaic virus, aphids, leafhoppers and grasshoppers fail to transmit the virus, 639
Basal stem lesions: of yellow birch seedlings, effect of fertilizer salt accumulation and low relative humidity on fungal invasion, 499
Bean: bacterial blight: virulent strain of Xanthomonas phaseoli from Uganda compared with strains from Nebraska and Colombia, 74; Phaseolus germplasm found tolerant to common blight bacterium, X. phaseoli, 111
---: Phaseolus aborigineus 283 and Costa Rica 1031 used to subdivide 7 races of Colletotrichum lindemuthianum found in Brazil within four race groups known in the U.S., Phaseolus vulgaris Cornell 49-242, was resistant to all races, 870
---: Phytophthora cryptogea isolated from bean in Kentucky, 190
---: Pythium and Sclerotinia rots of snap beans were controlled with a postharvest 30-sec dip in water at 125°C or with heated DCNA, 234
---: white mold disease of Great Northern bean, caused by Whetzelinia sclerotiorum, reduced with wider row spacing and determinate growth habit, 1070
Begonia: oxamyl and BAY 68138 controlled foliar nematodes in Rieger begonia, 603; foliar nematodes controlled effectively with oxamyl and aldicarb, 1015
Belonolaimus longicaudatus: 750
Bemisia tabaci: 463; 800
Beta vulgaris (see also Sugarbeet): sugarbeet cultivars introduced or grown in the Red River Valley were tested for resistance to Phoma storage rot, 204
Betula alleghaniensis: Diaporthe alleghaniensis cankering in yellow birch caused damage to 8-year-old released seedlings, 230; development of stem lesions on yellow birch seedlings grown in vermiculite caused by accumulation of fertilizer salts by evaporation, and low relative humidity, 499
Bioassay: local lesion response of Cassia occidentalis to cymbidium mosaic virus was not affected by treatment with benomyl to control powdery mildew, 475; small quantities of plant extract required for bioassay with chestnut blight fungus, 672
Biological control: of Stereum purpureum on plum tree wounds by inoculation with Trichoderma viride by spraying and also through pruning shears, 25; survival of sclerotia of Rhizoctonia solani in lake water, as possible biocontrol for water hyacinth, 601; strain of Colletotrichum gloeosporioides which controls northern jointvetch in rice fields is endemic in 31 rice-growing counties in Arkansas, 695; antagonism of Trichoderma spp. to sclerotia of Typhula incarnata, 924
Birch (see Betula alleghaniensis)
Birdsfoot trefoil: control of Pratylenchus penetrans with oxamyl, 237
Bitterbrush (Purshia tridentata): host of Fomes annosus, 407
Blackleg: of potato, seedpiece treatment with benomyl reduced incidence, 803
Black shank: of tobacco, race 1 of the pathogen found for first time in Georgia, 832
Blight: petal blight of chrysanthemum, caused by Itersonilia perplexans, in Florida, 181; Phomopsis blight, benomyl gave good control on eastern redcedar, 315; Rhizoctonia blight of grain sorghum foliage caused local epiphytotic in Georgia, 549; late blight of tomato, fungicidal control caused fruit damage, 612; seedling, of cotton, caused by three Pythium species, 848; Coryneum blight of peach in Iraq, 855; foliar blight of Santolina spp., caused by Alternaria tenuissima, 1072
---: Ascochyta: of chrysanthemum, caused by Mycosphaerella ligulicola, controlled

on unrooted cuttings with benomyl dip, 624; of chickpea, A. rabiei controlled with benomyl and thiabendazole, 742
---: bacterial: strains of Xanthomonas phaseoli, cause of common bean blight, from Uganda, Colombia, and Nebraska compared for virulence, 74; tolerance found in several cultivars and species of Phaseolus in tests in Nebraska, 111; field-overwintered Pseudomonas glycinea as possible primary inoculum of soybeans, 156; persistence in rice plant tissue, 227; of Gerbera jamesonii, caused by Pseudomonas cichorii, 504; of rice, leaf clipping technique for evaluating resistance to Xanthomonas oryzae, 537
---: Cercospora blight: controlled on eastern red cedar with benomyl, 315; controlled on arborvitae with benomyl sprays, 739
---: chestnut blight: control attempted with hypovirulent cultures of the pathogen, 225; micro-bioassay method applied to chestnut blight fungus, 672
---: early blight: of celery, Cercospora apii strains resistant to benomyl, 837; of tomato, caused by Alternaria solani, it is possible to select plants having resistance to both phases of early blight, 964
---: fire blight: of pear: for the last 6 years Egypt has been free of the disease, 646; a new inoculation device, and a simple humidity chamber described for screening large numbers of pear seedlings for resistance, 20
---: leaf: a "new" Helminthosporium leaf blight of corn found in the northern Corn Belt, 195; Helminthosporium blight of wheat, caused by H. sorokinianum, in India, 278; the role of temperature, leaf wetness and host age, in development of Alternaria triticina infection on wheat, 1000
---: ---: (northern corn leaf blight): of corn, caused by Helminthosporium turcicum, monogenic and multigenic resistance, 586; virulence of two H. turcicum populations was not changed following six successive passages through monogenic resistant or susceptible corn seedlings, 590
---: southern corn leaf blight (see Corn)
---: tip or shoot blight: of Pinus spp., caused by Sirococcus strobilinus, found for first time in the Western United States, in California, 69; S. strobilinus has been identified as the cause of a shoot blight of red pine in the Lake States, 246
Blight tolerance: to common blight bacterium of bean, tolerance to Xanthomonas phaseoli in several lines of Phaseolus species was always associated with late maturity, 111
Blossom symptoms: virus-induced symptoms on the blossom parts of apple, 359
Blueberry: anthracnose, gray mold rot and Alternaria rot made up 98% of all decays on packaged New Jersey blueberries immediately after harvest, after 4 days at 60°F, and in retail during 1970-72, 487; a survey to determine losses due to witches' broom of blueberry caused by rust fungus Pucciniastrum goeppertianum, in Maritime Provinces, 608; contributed to needle rust of balsam fir in Maritime Provinces, 765
Boll infection: of cotton, by Colletotrichum gossypii, 828
Boll rot (see Rot)
Botryosphaeria ribis: 28
Botrytis cinerea: 31; 67; 131; 210; 487; 760; 795; 969
Brazil: 870
British Columbia: 358; 359
British Solomon Islands: 701; 856
Broussonetia papyrifera (paper-mulberry): Corynespora cassiicola reported as cause of frog-eye lesions on leaves, disease was observed and collected by George Washington Carver in the 1930's, 296
Brown rot (see Rot)
Brown spot disease: of white field corn, partially attributed to change in tillage practices of not plowing under crop debris, 630
Brown spot needlecast (see Needlecast)
Brown stain: of onion, caused by Botrytis cinerea, 13 cultivars tested for incidence, severity and type of brown stain, in New York, 210
Buckwheat: a good host for Pratylenchus penetrans, 728
Bunchy top disease: of papaya, first report of leafhopper Empoasca stevensi as a vector, 412

Cabbage (see Crucifers)
Caladium: root-knot nematodes and other plant-parasitic nematodes associated with Caladium in Florida, 558
California: 44; 67; 69; 115; 192; 258; 619; 634; 719; 904; 926; 1066
Cambial death rate: in relation to invasion of Fomes annosus in infection wounds of trees treated with herbicides, 94
Cameroun: 529
Canada: 237
Canker: pitch canker, caused by Fusarium

lateritium f. pini found on eastern white pine in Virginia for the first time, 182; of yellow birch seedlings, by Diaporthe alleghaniensis near Dukes, Michigan, 230; Monochaetia canker of Arizona cypress, controlled with benomyl, 315; Scleroderris canker on red and Scots pine in Vermont, 338; canker and branch dieback of apple trees in Wisconsin caused by Stereum purpureum, 473; Apioporthe canker of filbert (Corylus) and Monochaetia canker of arborvitae found in western Washington, 522; black rot canker of apple, cultivar susceptibility, 676
Canker: bacterial: of tomato: high resistance in new tomato variety, H2990, 584; spread during clipping of transplants, 767; loss in a resistant and susceptible cultivar, 787; control by seed treatment, 974
Canker stain disease: of planetree, caused by Ceratocystis fimbriata f. platani, a tree wound dressing + benomyl prevented growth of the pathogen at wounding sites, 28
Carica papaya (see Papaya)
Carnation: bacterial wilt caused by Erwinia chrysanthemi is identified in New Zealand for the first time, 136
Carrot: postharvest decay, benomyl and DCNA compared favorably with SOPP as dips for control of postharvest decay, 697; effects of bacterial contamination and refrigerated storage on bacterial soft rot, 896
Carver, George Washington: a fungal pathogen collected by Carver in the 1930's on leaves of Broussonetia papyrifera (paper-mulberry) identified as Corynespora cassiicola, 296
Cassava (Manihot esculenta): bacterial wilt, caused by Xanthomonas manihotis, 824
Cassia occidentalis: benomyl completely controlled powdery mildew on this plant that is used as a local-lesion host for Cymbidium mosaic virus, 475
Celery (Apium graveolens): Septoria leaf spot, controlled with several nonsystemic fungicides and the systemic fungicide benomyl, 425; early blight, caused by Cercospora apii, which has naturally occurring benomyl-resistant strains, was controlled with chlorothalonil in Florida tests, 837
Central America: 47
Cephalosporium gramineum: 187
--- gregatum (= Phialophora gregata): 269; 660; 853; 1063
Ceratocystis fagacearum: 254; 325; 1055
--- fimbriata f. platani: 28
--- ulmi: 35; 511; 576; 701; 997
Cercospora sp.: 315
--- spp.: 770
--- apii: 837
--- beticola: 321
--- fusimaculans: 33
--- sequoiae: 739
--- --- var. juniperi: 315
Cercosporella herpotrichoides: 187; 935
Chalara quercina (Ceratocystis fagacearum): 325
Chemicals (see also Antibiotics, Fungicides and Nematicides): a device described that can inject chemicals into trees under pressure that is cheap, easy to build, and easy to install, 338; a new trunk injection technique for systemic chemicals, 513; hydrogen peroxide, for extraction of nematodes from banana roots, 678; indole-3-acetic acid (IAA), 2, 3, 5-triiodobenzoic acid (TIBA), and 6-furfurylaminopurine (kinetin) growth regulators decreased disease severity of charcoal rot of soybean, 789; lactophenol fixation method for nematodes, 1045
---: chlorination: chlorination of hydrocooling water for peaches and its effect on conidia of Monilinia fructicola, 814; sodium hypochlorite (NaOCl) controlled damping-off fungus of statice, 969; seed treatment with NaOCl reduced bacterial leaf and flower spot of zinnia, 1020; egg mass matrices of Meloidogyne spp. dissolved in NaOCl solution, 1025
---: herbicides, chemicals producing rapid cambial death, such as MSMA, are recommended for precommercial thinning of western hemlock, but not ones that produce slow senescence, as picloram-2, 4-D, 94
---: insecticides: azinphosmethyl, mixed with dodine is phytotoxic to cherry fruit, 428; monocrotophos, movement of carboxin in cotton seedling enhanced by coating of monocrotophos, 642; disulfoton, cowpeas planted inside a disulfoton-treated pearl millet trap-crop developed less cowpea mosaic virus than plants outside the trap-crop, 684; several tested together to control foliar nematode of Rieger begonia, 1015
---: liquid nitrogen: as storage medium for Macrophomina phaseolina sclerotia, 478; to preserve downy mildew (Phytophthora phaseoli) inoculum, 724; successful preservation of rust spores in, 793
Chemotherapeutic agents: two antibiotics and B. P. -101 reduced greening disease symptoms on lime plants, only B. P. -101 injected into stem brought about

complete recovery, 160
Cherry: pollen transmission of Prunus ringspot virus in 'Montmorency' cherry, 241; phytotoxicity of dodine and azinphosmethyl to cherry fruit, 428
Chestnut blight: use of hypovirulent cultures of Endothia parasitica from France as a possible protection against an American virulent strain of the pathogen in American chestnut trees, 225; bioassay method successfully tested in vitro growth of E. parasitica, but there is a lack of correlation with Castanea resistance patterns in nature, 672
Chickpea (Cicer arietinum): benomyl and thiabendazole were effective as seed-treatment fungicides for control of Ascochyta rabiei, 742
Christmas tree diseases: brown spot needlecast of Scotch pine, 55; benomyl gave good control of Cercospora and Phomopsis blights of eastern redcedar and Monochaetia canker of Arizona cypress in South Carolina tests, 315; needle rust of balsam fir, 765; Swiss needlecast found on Douglas-fir, 984
Chrysanthemum: Itersonilia petal blight found for first time in Florida, and first time since initial Minnesota report in 1956, 181; control of Ascochyta decay with 5-sec benomyl dip, 624
Citrus: lime, plants with greening symptoms when treated with B. P. -101, a new chemotherapeutic agent, injected into the stem, recovered and bore normal foliage and fruits, 160; unusual brown rot outbreak on grapefruit in Florida caused by Phytophthora-infested irrigation water, 391; P. parasitica resistance in seedlings, method for large scale screening, 669; postharvest decay of citrus fruits, benomyl, thiabendazole and thiophanate methyl provided effective control, 899
---: greasy spot: differential susceptibility of grapefruit on various rootstocks to Mycosphaerella citri, 294; benomyl and Difolatan controlled M. citri as well as copper fungicides, but Difolatan caused damage to fruit, 691
---: virus: first report of Satsuma dwarf virus disease on Satsuma mandarins in Turkey, 149; mechanical inoculation procedures for citrus exocortis virus, 886; Arizona 861 'Etrog' citron monoembryonic seedlings apparently can be used reliably as indicator plants for citrus exocortis virus, 1010
Cladosporium fulvum: 179; 612
--- herbarum: 756
Climatic factors: their influence on colonization of Helminthosporium maydis in corn hybrid following infection, 145; as they affect sporulation of an isolate of race T of H. maydis on a susceptible male-sterile corn hybrid, 304
Clipping: spread of bacterial canker of tomato transplants during clipping, 767
Cob parenchyma: because of uniformity of cells, this tissue may be ideal for study of host-pathogen interactions of corn stalk rot fungi such as Diplodia maydis, 308
Cocoa (Theobroma cacao): mating types were determined for 38 cultures of Phytophthora palmivora in Cameroun, 529
Coconut: no major parasitic diseases found in British Solomon Islands, leaf spots and blights were the most conspicuous diseases, 856
Coleosporium asterum: 793
Colletotrichum sp.: 747
--- dematium: 445
--- destructivum: 104
--- falcatum (see Physalospora tucumanensis)
--- gloeosporioides: 173
--- --- f. sp. aeschynomene: 695
--- gossypii: 756; 828
--- graminicola: 143; 272
--- lindemuthianum: 870
--- truncatum: 770
Colocasia esculenta (taro): corm rot, caused by Fusarium oxysporum, recorded for first time on Colocasia in the British Solomon Islands, 701
Colorado: 511
Conidia: of Lecanosticta acicola, the imperfect stage of brown spot needlecast, were the most important method of disease spread between closely planted Scotch pine Christmas trees, 55
Conifers (see also Species names): Rhizina root rot in conifer plantations in western Washington, 167; listing of location and tree species in the intermountain region of the U. S. where Fomes annosus has been found, 407
Control: of witches' broom of blueberry, eradication of the brooms, plus herbicide kill of plants infected with Exobasidium vaccinii should be considered, 608
Corn: downy mildew, caused by Sclerospora sorghi, several corn inbreds not reported previously show resistance, 42; severe epiphytotic of anthracnose occurred in 'Jubilee' sweet corn in Benton County, Indiana in mid-August 1972, 143; a Helminthosporium sp. has been identified as the cause of an apparently new disease of corn in the Corn Belt, 195; Goss's wilt (also called leaf freckles and wilt), its occurrence and spread in western Corn Belt from 1969-1972, 291;

Corn (continued): parenchyma cell death and Diplodia maydis susceptibility in stalks and ears of corn, 308; corn stunt, spiroplasmas occur naturally in corn stunt-infected plants in Mexico and support hypothesis that these organisms are the CS disease agent, 333; Physoderma maydis infection in Illinois on susceptible corn inbreds and hybrids planted in overwintered debris from infected fields in Illinois, 630; effect of ozone on 'Golden Jubilee' sweet corn, 719; a different Helminthosporium, "x" found in Connecticut and Iowa, mates readily with H. carbonum, and could be a new race, 780; new race of H. carbonum, 822; stalk rot pathogen, Macrophomina phaseoli, survives in corn and sorghum stalk residue as sclerotia for over a year, 873; a rotational crop for peanuts, corn is resistant to Cylindrocladium crotalariae, 1035; increases in host nuclear dry mass, nuclear area, and nucleolar area in Diplodia-infected corn, 1043

---: northern corn leaf blight: monogenic and polygenic forms of resistance in corn to Helminthosporium turcicum complement each other and both can be used together, 586; no virulence change occurred in two H. turcicum populations following serial host passage, 590

Corn: southern corn leaf blight (SCLB): further studies indicate that race T of Helminthosporium maydis has been the predominant member of the pathogen population for many years, 18; effect of duration of high humidity on lesion production by H. maydis on corn seedlings, 76; several climatic factors greatly affect the colonization of a susceptible corn hybrid by an isolate of race T of H. maydis, 145; isolates of race T of H. maydis collected before 1970 found to be pathogenic to corn hybrids with male-sterile cytoplasm, 253; the Tms factor increased the rate of blight development 2.5 times that of the N corn in a mixed stand, 276; in 1971 in seven States spore sampling of H. maydis race T with air-suction type samplers resulted in disappointingly low spore counts at all stations, 298; seed infected with H. maydis race T and treated with carboxin + thiram greatly increased numbers of surviving seedlings, 344; race types of H. maydis found on hybrids in normal cytoplasm in 1972 in Pennsylvania, 411; toxin activity of culture filtrates of six isolates of H. maydis from races O and T, and toxin specificity, 420; popcorn is susceptible to both H. maydis races O and T and to race 2 of H. carbonum, 507; evidence of race T of H. maydis in Sierra Leone, 509; susceptibility of detached corn leaves to H. maydis races O and T was determined by using disease ratings, electrolyte leakage and peroxidase leakage, 518; effect of relative humidity and air speed on release of H. maydis conidia, 627; three formulae developed to calculate quantitatively the sporulation rate of an isolate of H. maydis race T under a given set of environmental conditions, 730; cross-protection by race O against race T of H. maydis, 971; incidence of race O and T of H. maydis at Gainesville, Florida, 981; number of lesions on susceptible P-A-G 15029^{T} and resistant Pio. 3369A^{T} cultivars was not significantly different but lesion size and number of conidia were significantly reduced on resistant Pio. 3369A^{T}, 1047

---: virus: a mosaic virus disease of maize and sorghum in Israel is similar to maize dwarf mosaic virus of the U.S., 125; three mechanically transmissible viruses were isolated and identified from corn and sorghum in South Dakota in 1972, 373; maize streak disease, results support the report that MSV from Africa may be a small nuclear-infecting polyhedron, 414

Corn Belt: occurrence and spread of Goss's bacterial wilt of corn, in three States, 291

Corynebacterium sp: 291

---michiganense: 258; 584; 767; 787; 974

Corynespora cassiicola: 296; 375; 404; 921

Coryneum beijerinckii: 855

Cotton: seed treatment with systemic nematicides for root-knot nematode control was not effective, 129; an analysis of cotton disease losses from 1965 through 1971 indicates losses due to nematodes and boll rots have increased on a national basis, geographically, disease losses have increased in the West, 325; a 25-month study showed the population levels of Rotylenchulus reniformis increased about 45 days after seeding a new crop, 399; use of herbicides in cotton fields has resulted in increased populations of nutsedges that are hosts of plant-parasitic nematodes, 402; reduction in yield of cotton caused by parasitic diseases in 1972, 403; root drench and foliar applications of CW-524 at fairly high concentrations delayed disease symptoms on cotton following inoculation with Verticillium albo-atrum, 460; seedling protection by the fungicide carboxin

improved with monocrotophos insecticide, 642; Aspergillus flavus boll rot infection in the U. S. Cotton Belt in relation to high aqueous-extract pH of fiber, 664; microorganisms associated with cotton fiber samples from across the U. S., 756; geographical localization of cotton boll infections with Colletotrichum gossypii, 828; Pythium sylvaticum, P. ultimum, and P. irregulare identified as seedling blight pathogens, 848; level of microbial infection of cotton seeds in the field with respect to pH of aqueous extracts of fiber, 994; peanuts following cotton and tobacco in rotation were damaged by Cylindrocladium crotalariae in North Carolina, 1035

Cottony leak: of bean, caused by Pythium butleri, controlled in simulated transit and storage trial with hot water and DCNA, 234; of ripe tomato fruit, caused by Pythium spp., 1066

Cover crops: oats, sudex, and sudangrass were least favorable hosts for Pratylenchus penetrans of seven cover crop hosts tested, 728

Cowpea (Vigna sinensis): inoculation studies show that the cowpea strain of southern bean mosaic virus is seed-transmitted in two varieties of cowpea, 13; cowpea plants growing within an insecticide-treated pearl millet trap-crop had less virus infection than the nonscreened control plants, 684; Hoplolaimus seinhorsti is endoparasitic in cowpea roots in Nigeria, 798

Crambe spp.: world listing with citations, of parasitic fungi, bacteria and viruses that have been reported on species of Crambe, 704

Criconemoides sp.: 891

--- xenoplax: 115

Cronartium comandrae: 17; 283

--- fusiforme: 776; 793; 1013

Crown class: effect of crown class on canker healing and tree mortality in yellow birch seedlings exposed to Diaporthe alleghaniensis, 230

Crownvetch (Coronilla varia): Colletotrichum destructivum is cause of anthracnose of crownvetch in Pennsylvania, 104

Crucifers: cabbage, aerial application of benomyl was very effective in control of Sclerotinia sclerotiorum in cabbage seed fields in Washington, 164; turnip and mustard carry a low infection of turnip mosaic virus and cauliflower mosaic virus in Georgia, 978

Cucumis melo (see Cucurbits)

Cucurbits: yellow summer squash, chlorotic leaf spotting caused by the severe strain of bean yellow mosaic virus, 280; Cucumis melo, bacilliform particles in cortex cells of Cucumis melo virus-infected fruits, in Spain, 649; cucumber, nematodes controlled in greenhouse with methyl bromide and DD, 1032

Culture medium: effect of culture medium on production and pathogenicity of Alternaria alternata conidia, 39

Cupressus arizonica (Arizona cypress): benomyl controlled Phomopsis blight in a nursery, 59; benomyl was best of several fungicides tested for control of Monochaetia canker in Christmas tree study, 315

Cutting decay: of geranium, caused by Fusarium and Rhizoctonia, 177; of Gynura aurantiaca and Hoya carnosa caused by Rhizoctonia solani, controlled with benomyl, 222; of chrysanthemum cuttings, controlled with benomyl, 624

Cylindrocladium crotalariae: 90; 387; 1035

--- scoparium: 500

Cyperus esculentus (see Nutsedge)

Cytoplasm (male-sterile): of corn: effect of high humidity on lesion development of Helminthosporium maydis on corn seedlings with male-sterile or normal cytoplasm, 76; climatic factors affect colonization of susceptible corn hybrid by H. maydis race T, 145; H. maydis race T isolates collected before 1970 are pathogenic to male-sterile corn hybrids, 253; progress of SCLB in a mixed stand of normal and Tms cytoplasm corn, 276; male-sterile corn seed infected with H. maydis race T had higher stand counts after seed treatment with carboxin + thiram, 344; pathogenicity of various isolates of H. maydis from races O and T on Fla-200A (T-cytoplasm) in relation to toxin activity of culture filtrates on the same corn line, 420; H. maydis race T in Sierra Leone, 509; regression equations for predicting sporulation of an isolate of race T of H. maydis on a susceptible male-sterile corn hybrid, 730; cross-protection of race O and T of H. maydis, 971; predominance of race T of H. maydis in corn containing male-sterile cytoplasm and race O on normal cytoplasm corn in Florida 1971-1972, 981; effect of some resistance on the development of SCLB in two male-sterile hybrids, 1047

Cytospora cincta: 28

Dactynotus escalantii: first report as vector of pea seedborne mosaic virus, 552

Damping-off: of pea, Pythium ultimum, controlled with thiram and chloroneb

combination, 3; of statice, Botrytis cinerea controlled with hot water, benomyl, or Clorox soaks, 969
Datura metel var. muricata: Alternaria crassa found to be a seedborne fungus in Datura, 960
Decline: of apple in Wisconsin, Stereum purpureum was associated with cankers and branch dieback, 473; of bananas in Lebanon, 615; of pear, controlled in California with oxytetracycline hydrochloride treatments, 634
Delaware: 542
Determinate growth: in Great Northern beans helped to reduce incidence of white mold disease, caused by Whetzelinia sclerotiorum, 1070
Diaporthe alleghaniensis: 230
Dieback: Sirococcus tip dieback of Pinus spp. in California forest nurseries, 69; of filbert, caused by Apioporthe anomala, found in Washington, 522
Diplodia maydis: 308; 1043
Dips: control of postharvest decay of stone fruits with hot water and fungicides, 62; benomyl dips at 55°C controlled postharvest decay of mango, 173; control of Pythium and Sclerotinia rots of snap beans with postharvest hot water and DCNA dips, 234; bare-root dips of thionazin and BAY 25141 completely controlled root-knot-infected peach trees, 489; 15-minute hot water dip controlled Pratylenchus penetrans in raspberry root cuttings, 492; control of Ascochyta decay of chrysanthemum cuttings with benomyl, 624; fungicide dips for control of decay of carrots in storage, 697
Diseases: of Crambe spp., 704
Distribution: distribution, frequency of occurrence and population dynamics of Rotylenchulus reniformis in Georgia, 399; known distribution of Fomes annosus in the intermountain region of the U. S., 407; of mating types of Phytophthora palmivora in Cameroun, 529; distribution and factors influencing distribution of Douglas-fir dwarf mistletoe, 716; of chlorotic leaf spot virus in apple budsticks, 865; of pepper viruses in Florida and their identification, 991
Ditylenchus radicicola: 1021
Douglas-fir (see Pseudotsuga menziesii)
Downy mildew: Stenotaphrum secundatum on St. Augustinegrass, found for the first time, in Louisiana, 16; of corn, an additional series of corn inbreds were tested for their reaction to Sclerospora sorghi, 42; Phytophthora phaseoli infection on lima bean epicotyl stored in liquid nitrogen was virulent after 12 months' storage, 724
Drechslera gigantea: 84
Dutch elm disease (DED): long-term protective action of benomyl soil treatment against DED, 35; locations in Colorado in 1972, 511; determination of Ceratocystis ulmi sexual types present in Wyoming, 576; first report of the disease in Montana, 701; resistance in Chinese elm hybrids, 997
Dwarf mistletoe: on lodgepole pine in Montana, a new distribution record, 298; additional sites of dwarf mistletoe west of the Cascade Divide, 716; detection on jack pine with aerial photographs, 941

Eggplant (Solanum melongena): is alternate host for Puccinia substriata var. indica that causes rust on pearl millet and can serve as reservoir for the disease, 262; no resistance found in cultivars or breeding material of S. melongena to root-knot nematode, Meloidogyne incognita, 606
Egypt: 646
Electrolyte and peroxidase leakage: as indicators of susceptibility of maize inbreds to Helminthosporium maydis races O and T, 518
Electron microscopy: of bean yellow mosaic virus, 280; of Hydrangea ringspot virus, 300
Elm: American: phloem necrosis suppressed on American and cedar elms with injected tetracycline antibiotics, 341; evaluation of new trunk injection technique in American elms with systemic chemicals, 513
---: Chinese, resistance to DED, 997
---: Dutch elm disease (DED): 35; 511; 576; 701; 997
Empoasca stevensi: 412
Endothia parasitica: 225; 672
Epicoccum cocos: 856
Epidemic: of wheat stem rust in 1972 was averted by use of early maturing and resistant varieties and nearly complete eradication of barberries, 434
Epidemiological study: moisture modification shelters designed for epidemiological studies of foliar diseases, 329
Epidemiology: of stem, leaf and stripe rusts of wheat in northern India, 8; Sirococcus strobilinus development on Pinus spp. in California appears to be restricted by temperatures below 10°C, 69; of Lophodermium pinastri and control of needlecast of Scotch pine in Ohio with maneb, 515; development of leaf blight of wheat, caused by Alternaria triticina, 1000
Epiphytotic: of anthracnose, caused by Colletotrichum graminicola, occurred

in Benton County, Indiana on 'Jubilee' sweet corn, 143; local epiphytotic occurred on grain sorghum in research plots, by Rhizoctonia solani, 549; a severe epiphytotic of brown spot disease on certain lines of white field corn in Illinois in 1971, 630
Erwinia sp.: 453
--- spp.: 803
--- amylovora: 646
--- carotovora: 896
--- chrysanthemi: 136
Erysiphe graminis f. sp. tritici: 187
Ethylene gas: as a cause of injury to Easter lilies, 1023
Euonymus: Mycocentrospora sp. identified as cause of an apparently new leaf spot disease of Euonymus in Ohio, 807
Eyespot: of wheat, 187

False toadflax (Comandra umbellata ssp. umbellata): the alternate host for comandra blister rust was present in Kentucky near a loblolly plantation found infected for the first time with Cronartium comandrae, 17
Felling: felling oak wilt-diseased trees did not prevent bark beetle breeding but was more effective in reducing beetle density and fungus mat formation than girdling, 1055
Fertility level: of St. Augustinegrass, the lower the level of fertility, the higher the incidence of leaf spot, caused by Cercospora fusimaculans, 33
Fertilizer: applying fertilizer to brush piles of apple prunings or covering with plastic sheeting reduced sporulation of Physalospora obtusa by 75 to 90%, 158; effect of fertilizer accumulation and low relative humidity on development of stem lesions on yellow birch seedlings, 499; high fertilizer-high mowing of Kentucky bluegrass in a majority of lines caused greater leaf spot infection than low fertilizer-low mowing, 544
Fire blight (see Blight)
Flindersia brayleyana (Queensland maple): seedling disease caused by Rhizoctonia solani was controlled with seed coatings of PCNB, 907
Florida: 119; 122; 181; 260; 391; 558; 612; 638; 691; 747; 837; 861; 981; 991
Foliar diseases: moisture modification shelters tested for epidemiological studies of foliar diseases, such as peanut leafspot, 329; of alfalfa, controlled with zinc + maneb sprays in Minnesota and yield of hay was increased, 353
Fomes annosus: 94; 407
--- nigrolimitatus: 346
Fraxinus americana (white ash): sequential development of viruslike symptoms in ash over a 6-year period, 396; white ash and green ash (F. pennsylvanica), response to chronic ozone fumigation, 914
Fruit decay: (see Rot)
Fruit rot (see Rot)
Fungi: decline of bananas in Lebanon thought to result from interaction between nematodes and Fusarium oxysporum and/or Rhizoctonia solani, 615; fungi, bacteria and viruses reported on Crambe spp., a world listing, 704; fungi infecting cotton fiber samples in relation to geographical location of growth, 756
Fungicides (see also Antibiotics, Chemicals, and Nematicides): several tested together: thiophanate methyl, thiabendazole, chlorothalonil, Dyrene, and BAS 3460-F all controlled Cercospora fusimaculans on St. Augustinegrass, 33; heat and fungicide treatments to control postharvest brown rot of stone fruits, 62, benomyl was best of 4 fungicides tested for control of Botrytis storage rot of giant sequoia seedlings, 67; some metallic compounds other than arsenic salts will control ball moss, that grows on the bark of some Texas shade trees, 81; Difolatan, NIA 5961, and PCNB-ETMT controlled Cylindrocladium black rot on peanuts for 65 days after planting, 90; for control of Christmas tree diseases, benomyl was best of six tested for control of Cercospora and Phomopsis blights, 315; as seed treatment for corn infected with Helminthosporium maydis race T, carboxin + thiram proved a beneficial treatment for improved seedling stand, 344; zinc + maneb, Polyram, Dyrene, chlorothalonil, and benomyl gave good control of Septoria leafspot of celery, 425; none improved the germination of snapdragon seed heavily infested with Alternaria alternata, Phyllosticta antirrhini, or Stemphylium botryosum, 592; chlorothalonil, Polyram, zinc ion + maneb, and zinc ion + maneb + Dyrene controlled late blight of tomato, 612; benomyl and Difolatan compared favorably with copper fungicides for control of greasy spot of citrus, 691; Dyrene, carboxin and BAS 3191 F inhibited growth of puffball fairy ring fungus in bentgrass turf, 711; benomyl and thiabendazole reduced incidence of Ascochyta blight in chickpea seedlings, 742; where smut was not present, no seed treatments improved plant stand or yield of 'Nugaines' winter wheat, 1006
---: acetaldehyde, for control of Botrytis

Fungicides (continued): cinerea on raspberry, 795
---: benomyl: added to an asphaltum-based wound dressing, was effective in prevention of growth of Ceratocystis fimbriata f. platani, Verticillium dahliae, Botryosphaeria ribis, and Cytospora cincta on wounds on planetree, maple, apple and peach with no adverse effect on callus formation, 28; provided excellent control of gray mold of strawberry, in Poland, 31; pressure-injected into soil, benomyl protected elm trees inoculated with Ceratocystis ulmi from foliar symptoms for three consecutive summers, 35; curative action of foliar sprays against Fusarium wilt of tomato, 37; controlled Phomopsis blight of nursery seedlings of Arizona cypress, 59; benomyl as an aerial spray and cyanamide as a ground application, protected cabbage seed fields against Sclerotinia sclerotiorum, 164; controlled postharvest decay of mango in South Africa, 173; effective against cutting decay on Gynura aurantiaca and Hoya carnosa, 222; controlled Lophodermium pinastri on red pine seedlings, 263; prevented disease development in Fusarium-inoculated tomato plants when added to soil pasteurized with DD-MENCS, 267; development in Greece of benomyl-resistant strains of Cercospora beticola, cause of leaf spot of sugarbeet, 321; as a soil drench controlled wilt of Pyracantha caused by Fusarium oxysporum, 389; was better than thiophanate methyl or thiabendazole for control of tomato fruit rots, 453; gave complete control of powdery mildew of Cassia occidentalis, 475; benomyl and chlorothalonil gave good control as protective sprays against Cylindrocladium scoparium infection of silver buttonwood cuttings, 500; technique for trunk injection with benomyl, 513; efficacy of benomyl spray for control of apple scab was enhanced by addition of 1% summer superior oil in South Australia, 560; as 5-sec dip, controlled Ascochyta blight of chrysanthemum unrooted cuttings, 624; reduced postharvest decay in carrots, 697; sprays at 2- to 4-week intervals controlled arborvitae blight, 739; reduced Rhizoctonia and black leg on potato, 803; disease progress of Cercospora apii resistant to benomyl, 837; benomyl, thiabendazole and thiophanate methyl in low concentrations controlled postharvest decay in citrus, 899; gave control of Sclerotinia sclerotiorum in tobacco seedbeds, as drench, 932; no benomyl tolerance found in Cercosporella herpotrichoides, the cause of strawbreaker foot rot of wheat, 935; controlled damping-off of statice, 969; ineffective against foliar nematode of Rieger begonia, 1015
---: benzyl isothiocyanate (BITC), a natural constituent of papaya, when used as dip or as a vapor reduced incidence of stem-end rot and Rhizopus stolonifer soft rot on green papaya fruit, not effective on mature fruit, 86
---: B. P. -101, as a curative of greening disease of citrus, 160
---: BT (4-n-butyl-1, 2, 4-triazole): a single application of BT in soil through irrigation water or on the crop as a spray controlled leaf rust of wheat, 572; single spray applications controlled wheat leaf rust, 653
---: carboxin: treating wheat planting seed with carboxin reduced incidence of smut infection in the seed crop produced, 187; gave excellent control of stripe rust on three cultivars of Kentucky bluegrass in Hawaii, 217; movement in cotton seedlings apparently improved with addition of insecticide monocrotophos, 642
---: Cela W 524: gave outstanding control of brown rot on plum, peach and tart cherry, 457; delayed disease symptoms of Verticillium wilt in cotton inoculated with the wilt fungus, 460; partially effective as soil drench against fusiform gall rust of slash pine, 776
---: chlorine, unheated dips of chlorine, SOPP, dehydroacetic acid were not effective against Pythium and Sclerotinia rots of snap beans, 234
---: chloroneb: synergism between chloroneb and thiram on pea seeds controlled seed rot and damping-off by Pythium ultimum, 3; was nonphytotoxic as spray but was phytotoxic as a drench for foliage plant cuttings, 222
---: chlorothalonil: controlled needlecast fungus, Lophodermium pinastri, on red pine seedlings, 263; gave good control of Cylindrocladium-blighted cuttings of silver buttonwood, 500; benomyl-resistant strain of Cercospora apii on celery controlled with chlorothalonil and was better than zinc + maneb sprays, 837
---: CO_2, California strawberries held in CO_2-enriched atmospheres had significantly less softening and decay than fruit held in air, 44
---: copper: K 101, controlled ball moss on shade trees in Texas, 81; copper fungicides are the most effective of fungicides recommended in Florida for citrus greasy spot, 691; tribasic copper sulfate, did not control spread of bacterial canker

Fungicides (continued): of tomato in diseased transplants, 767; copper compounds suppress Aphanomyces root rot of peas, 876
---: cyanamide, as a ground application prevented production of inoculum of Sclerotinia sclerotiorum in cabbage seed fields, 164
—-: cycloheximide, controlled needlecast fungus, Lophodermium pinastri, on red pine seedlings, 263
---: DCNA: in combination with benomyl in hot water as a dip was more effective than fungicidal dips in cold water, 62; a 30-sec heated dip controlled Pythium and Sclerotinia rots of snap beans when applied postharvest, 234; reduced storage rots on carrots in storage for processing, 697
---: DD-MENCS, tomato plants inoculated with Fusarium oxysporum f. sp. lycopersici and planted in soil treated with DD-MENCS developed more wilt than those planted in unpasteurized soil, 267
—-: dichlofluanid (Euparen), controlled Botrytis cinerea on strawberries and yields were increased, 31
---: dichlozoline, provided excellent protection against Botrytis cinerea in strawberry, 31
---: Difolatan: and cupric hydroxide not effective at rates tested for control of Phomopsis blight of Arizona cypress, 59; Difolatan alone was not phytotoxic to cherry fruit, but when mixed with azinphosmethyl was phytotoxic, 428
---: dinitro, was ineffective for control of fungi on wounds of planetree, maple, peach and apple when added to tree wound dressing, 28
---: DMTT, Sclerotinia in tobacco seedbeds was controlled by fumigation with DMTT or methyl bromide, 932
---: dodine: phytotoxic to cherry fruit with increasing concentrations and temperature above 70°F, 428; control of arborvitae blight, 739; tolerance of Venturia inaequalis in relation to usage of dodine in apple orchards, 817
---: HCl, Corynebacterium michiganense was eradicated from infected tomato seed by acid extraction followed by 3-hour drying period, 974
---: maneb: effective against needlecast fungus of red pine, 263; best of six fungicides tested for control of Lophodermium needlecast of Scotch pine, 515; reduced black leg of potato, 803
---: methyl bromide: at 400 lb/A under cover shows promise as preplant treatment against Armillaria mellea and nematodes in vineyard replants, 115; fumigation controlled Sclerotinia in tobacco seedbeds, 932
---: oxycarboxin: gave excellent control of stripe rust on Kentucky bluegrass, 217; as seed treatment gave initial control of leaf and stem rust of wheat, but dropped off after 70 to 83 days, 567; effective as soil drench against fusiform gall rust of slash pine, 776
---: PCNB: PCNB, chloroneb and ETMT + thiabendazole were less effective than benomyl for control of cutting decay of Gynura and Hoya, 222; tested for reduction of symptoms of Rhizoctonia solani and bacterial black leg, of potato, 803; as seed coating controlled Rhizoctonia disease of Queensland maple seedlings, 907
---: thiabendazole: as an additive to tree wound dressing for control of Ceratocystis fimbriata platani and other organisms, 28; gave satisfactory control of arborvitae blight, 739; used to control postharvest decay of citrus, 899
---: thiophanate methyl: excellent control of gray mold on strawberry, and an increase in berry yields, 31; as heated dip reduced Monilinia decay of stone fruits, 62; effective as fungicide for control of brown rot of stone fruits, 457; was as good as maneb in protecting Scotch pine against Lophodermium pinastri but was not better, 515; applied 47 days preharvest provided effective decay control of 'Marrs Early' oranges, 899; effective for control of Sclerotinia in tobacco seedbeds, 932
---: thiram: thiram and chloroneb combination better than either fungicide alone for control of seed rot and damping-off of pea seeds, 3; gave excellent control of Alternaria crassa on Datura seeds, 960
---: TPTH: gave good control of Lophodermium needlecast, 515; was as effective as benomyl in control of pecan scab on 'Mahan' pecan trees, 902
---: zinc + maneb: ineffective against several organisms when added to tree wound dressing, 28; effective in controlling foliar diseases of alfalfa, 353
---: ziram, not effective against needlecast fungus of red pine, 263
Fusarium sp.: 438
--- spp.: 177; 187; 403; 756; 770; 912
--- lateritium f. pini: 182
--- moniliforme: 318
--- oxysporum: 389; 615; 701; 1058
--- --- f. sp. gladioli: 957
--- --- f. sp. lycopersici: 37; 119; 267; 724
--- --- f. sp. vasinfectum: 403

--- roseum: 891; 1058
--- solani: 1058
--- tricinctum: 891
Fusicladium effusum: 902

Gall: basal gall of sunflowers, incited by Plasmopara halstedii, in North Dakota and Minnesota, 647
Geographic distribution: study on past occurrence and geographical distribution of isolates of race T of Helminthosporium maydis indicate that race T has been the predominant member of the pathogen population for many years, 18
Geographical location: in relation to prevalence of infecting organisms on cotton fiber, 756; localization of infections of cotton bolls with Colletotrichum gossypii, 828
Geographical spread: of Goss's bacterial wilt (leaf freckles and wilt) of corn in Nebraska, Kansas and Iowa over the last 4 years, 291
Georgia: 90; 399; 402; 549; 750; 832; 978
Geranium (see Pelargonium)
Gerbera jamesonii: severe leaf spot, caused by Pseudomonas cichorii, in Florida, 504
Germplasm: of Phaseolus spp., is tolerant of common blight bacterium, Xanthomonas phaseoli, 111
Gladiolus: Fusarium corm rot, six Fusarium isolates were single-spored and tested for virulence on five gladiolus cultivars, 957
Gloeosporium sp: 487
Glume blotch: of rice, caused by Pithomyces chartarum in India, 436
Gomphrena globosa (globe amaranth): seed transmission of tomato ringspot virus, 433
Gossypium (see Cotton)
Grand fir (see Abies grandis)
Grape: control of nematodes and Armillaria mellea in replants with methyl bromide, 115; Dogridge and Saltcreek cultivars were highly tolerant of nematode infection; Grenache, even with preplant fumigation did not establish itself, 416; grape-replant soil fumigation trials with 1, 3-D and methyl bromide in California gave excellent control of root-knot and other nematodes, 619
Graphium ulmi (Ceratocystis ulmi): 701
Grasses: several tested together: as hosts of Ditylenchus radicicola in Oregon hop yards, 1021; Brachiaria piligera, Imperata cylindrica var. major, and Paspalum conjugatum found as natural hosts of Xanthomonas albilineans in Queensland, 1040
---: bentgrass: a patch disease caused by Drechslera gigantea found on Kingston and Exeter cultivars in Rhode Island, 84; inhibition of puffball growth on bentgrass turf by fungicides, 711
---: bluegrass: Puccinia striiformis is reported for the first time from Hawaii on certain Kentucky bluegrass cultivars, 215; stripe rust controlled on three cultivars in Hawaii with carboxin and oxycarboxin, 217; low mowing-low fertilizer treatment caused less severe infection by Helminthosporium dictyoides than high mowing-high fertilizer treatment, 544; a complete range of susceptibility to Puccinia graminis found in 57 Poa pratensis introductions, an experimental cultivar and 4 standard cultivars tested, 595; stylet nematode genera and Fusarium species isolated from 'Merion' Kentucky bluegrass sod production fields in Pennsylvania, 891
---: johnsongrass, tested with several other grass species and found susceptible to Helminthosporium maydis races O and T and race 2 of H. carbonum, 507
---: Lolium-Festuca hybrids, selecting for resistance to Helminthosporium dictyoides, 563
---: ryegrass, Puccinia coronata found on perennial ryegrass cultivars in Hawaii, 215
---: sudangrass, was the least favorable host for Pratylenchus penetrans of seven cover crop species tested, 728
---: sudex, is poor host for Pratylenchus penetrans; as a cover crop it may help reduce populations of this nematode, 728
---: St. Augustinegrass: downy mildew and rust found on this grass in Louisiana, 16; susceptibility to Cercospora fusimaculans increases under low fertility regimes, almost complete control can be attained with fungicides, 33
Grasshopper: not a vector for barley stripe mosaic virus, 639
Gray mold: of strawberry, benomyl, dichlofluanid and thiophanate methyl gave excellent control in Poland, 31; control of Botrytis cinerea in stored giant sequoia seedlings with benomyl, 67; of blueberries in marketing channels, 487
Greasy spot: of citrus: disease severity on 'Marsh' grapefruit was influenced by the rootstock used, 294; benomyl and Difolatan gave control similar to copper fungicides; thiophanate methyl and chlorothalonil gave promise, 691
Greece: 321; 1032
Greening disease: of citrus, a new chemotherapeutic agent B. P. -101 was more effective than two antibiotics in curing

lime plants of greening symptoms, 160
Growth regulators: effective in reduction of charcoal rot of soybean, 789
Gymnosporangium spp.: 793
Gynura aurantiaca: Rhizoctonia solani cutting decay controlled effectively with benomyl, 222; studies on the extraction and mechanical transmission of citrus exocortis pathogen in gynura, 598

Haiti: 921
Hawaii: 86; 215; 217; 272; 907; 966
Helianthus annuus (see Sunflower)
Helicotylenchus multicinctus: 615
Helminthosporium sp.: a "new" Helminthosporium identified in Corn Belt, 195
--- "x": 780
--- spp.: 187
--- carbonum: 507; 822
--- dictyoides: 544; 563
--- gramineum: 495
--- maydis: 18; 76; 145; 253; 276; 298; 304; 344; 411; 420; 507; 509; 518; 627; 730; 971; 981; 1047
--- sorokinianum: 278
--- turcicum: 586; 590
Hemlock (see Tsuga heterophylla)
Heterodera latipons: 53
--- mothi: 946
--- schachtii: 942
Hibiscus cannabinus (see Kenaf)
--- sabdariffa (see Roselle)
Honduras: 47
Hoplolaimus columbus: 402
--- seinhorsti: 798
Hordeum leporinum: Septoria halophila, not previously recorded in Western Australia was observed on H. leporinum during the 1968 growing season, 367
Horizontal resistance: tolerance to race 2 of Fusarium oxysporum f. sp. lycopersici in tomato 'Homestead 24' was of little value in controlling Fusarium wilt in Florida test, 119; segregation for horizontal resistance to tomato early blight; factors for resistance to leaf spotting and stem canker segregate independently, 964
Host range: of severe strain of bean yellow mosaic virus, 280; of quail pea mosaic virus, 311; plants susceptible to the annual stock isolate of turnip mosaic virus found in Argentina, 379; for sting nematode, Belonolaimus longicaudatus, 750; of broad bean mottle virus, 833; of cucumber mosaic virus from Nandina domestica, 917; of Phytophthora cinnamomi in the Western Cape Province of South Africa, 1005; of Ditylenchus radicicola and its distribution in Willamette Valley, Oregon, 1021
Hot water (see Temperature)
Hoya carnosa ('Compacta'): cutting decay, caused by Rhizoctonia solani, controlled with benomyl, 222
Humidity: from 4 to 8-hr exposure to high humidity in mist chamber greatly increases lesion production by Helminthosporium maydis on corn seedlings, 76
Hydrangea: hydrangea ringspot virus is cause of phyllody of florists' Hydrangea, 300
Hypovirulent cultures: of Endothia parasitica, as a possible protectant against a virulent strain of the pathogen in American chestnut trees, 225

Idaho: 346; 803; 948
Illinois: 195; 630; 853
India: 8; 141; 160; 220; 278; 361; 436; 463; 495; 531; 572; 770
Indiana: 143
Inoculation: of two varieties of cowpea at different growth stages with southern bean mosaic virus, 13; of scarlet oak with gray and albino strains of Ceratocystis fagacearum alone and in paired combinations, 254; a new pressurized injector for forcing systemic chemicals into trees, 338; of elms infected with elm phloem necrosis, tetracycline antibiotics suppressed symptoms, 341; of sugarcane plants with Xanthomonas albilineans to screen for leaf scald resistance, 582; of rice with Xanthomonas oryzae, to study movement and multiplication in resistant and susceptible hosts, 784; of tomato varieties with Corynebacterium michiganense, to compare yield loss in resistant and susceptible hosts, 787; spray inoculation of pine seedlings with fusiform rust basidiospores is effective method for mass inoculation, 1013
Inoculum: optimum inoculum levels for screening alfalfa seedlings for resistance to the northern root-knot nematode, 657; comparison of methods of collecting inocula of root-knot nematode, 1025
Intermountain Region: the known distribution of Fomes annosus in this area, 407
Iowa: 471; 660; 780
Iraq: 855
Irradiation: effect of irradiation on growth and asexual reproduction of Botrytis cinerea, 131; gamma irradiation of mango fruits after hot dip treatment with benomyl did not noticeably improve decay control over benomyl alone, 173
Isolates: of Phialophora gregata (= Cephalosporium gregatum), 1063; 150 of 165 isolates of Helminthosporium maydis obtained from original specimens collected

from 1958-1967 throughout the world were race T and 15 were race O, 18
Isolation: of Phytophthora cryptogea from bean in Kentucky, the first report of an A2 mating type of Phytophthora in the State, 190
Israel: 53; 125
Itersonilia perplexans: 181

Japanese millet (Echinochloa crusgalli var. frumentacea): host of Pratylenchus penetrans, 728
Juniperus virginiana (eastern redcedar): benomyl was best of eight fungicides for control of Cercospora and Phomopsis blights on 2-year-old redcedar seedlings, 315

Kansas: 52; 291
Kenaf (Hibiscus cannabinus): resistance to Phytophthora in kenaf, roselle and other Hibiscus species, 884
Kentucky: 17; 190; 841; 845

Leaf curl: of apricot, caused by Taphrina deformans in Kulu Valley, India, 361
Leafhopper: not found to be vector for barley stripe mosaic virus, 639
Leaf mold: of tomato: one cultivar was resistant to Cladosporium fulvum, five showed moderate resistance, and all others tested ranged from moderate-severe to very severe in disease ratings, 179; fungicidal control caused fruit damage, 612
Leaf pucker disease: of apple, there appears to be an inverse correlation between symptom severity ratings and heat unit totals during the 40-day full bloom and post-bloom period, 358
Leaf scald: of rice, caused by Rhynchosporium oryzae, found in Louisiana, 715
---: of sugarcane: an inoculation technique for screening varieties for resistance to leaf scald, 582; naturally occurring grass hosts in Australia, 1040
Leaf spot: of St. Augustinegrass, caused by Cercospora fusimaculans, 33; Cercospora leaf spot, of sugarbeet, strains of C. beticola are present in Greece that are resistant to benomyl, 321; Corynespora leaf spot, a new disease of tobacco, identified in Nigeria, 375; of celery, caused by Septoria apiicola, 425; of Medicago spp., 58 accessions and five standard cultivars were evaluated for field resistance to Pseudopeziza medicaginis, 468; frogeye leaf spot of apple, caused by Physalospora obtusa, 676; Colletotrichum leaf spot and stem canker on Stylosanthes humilis, 747; of Euonymus, by Mycocentrospora sp., 807; new alfalfa variety, Apalachee, displayed high level of resistance to Stemphylium botryosum, 946;
---: bacterial: of zinnia, caused by Xanthomonas nigromaculans f. sp. zinniae, 555; bacterial leaf and flower spot of zinnia, 1020
Leaf stripe: of wheat, caused by Cephalosporium gramineum, 187
Leaf wetness: of wheat and its role in the development of leaf blight of wheat, 1000
Lebanon: 615
Lecanosticta acicola: 55
Lethal yellowing: of palms in Florida, found in Christmas palm (Veitchia merrilli) and fan palm (Pritchardia pacifica) in addition to coconut (Cocos nucifera), 861
Light: near-ultraviolet radiation was a limiting factor in the asexual sporulation of Botrytis cinerea; quality of radiation is an important factor in the sporulation of this pathogen, 131; its importance in the sporulation of Helminthosporium maydis and conidiophore formation, 304
Light microscopy: for the diagnosis of ratoon stunting disease of sugarcane, 987
Lily: Easter lily, injury caused by ethylene gas, 1023
Longidorus cohni: 53
Lophodermium pinastri: 263; 515
Louisiana: 16; 199; 420; 715; 811
Lyophilization: a promising method for preservation of barley yellow dwarf virus over long periods, 862

Macrophomina phaseoli (now Macrophomina phaseolina): 770; 873
--- phaseolina (= Macrophomina phaseoli): 318; 478; 770; 789; 873
Macrosiphum euphorbiae: 280; 552; 841
--- rosae: 552
Maize (see Corn)
Male-sterile cytoplasm (see Cytoplasm)
Mango (Mangifera indica): postharvest decay caused by anthracnose controlled in South Africa with benomyl dip, 173
Maple (Acer platanoides): tree wound dressing amended with fungicide controlled Verticillium dahliae infection, 28; A. rubrum, A. saccharinum, and A. saccharum seedlings tested for ozone tolerance; silver maple and sugar maple showed reductions in growth after exposure to ozone, 914
Maritime Provinces: 608; 765
Market diseases: of New Jersey blueberries, anthracnose, gray mold rot and Alternaria, 487
Massachusetts: 179
Mating types: of Phytophthora palmivora,

their distribution in Cameroun, 529
Matthiola incana (see Stock)
Medicago spp. (see also Alfalfa): field resistance to Pseudopeziza medicaginis in alfalfa accessions and cultivars, 468
Melampsora spp.: 793
Meloidogyne sp.: 403
--- spp.: 558; 1025
--- hapla: 505; 657
--- incognita: 115; 199; 371; 402; 416; 489; 606; 615; 921; 1032; 1033
--- --- acrita: 129
--- javanica: 141
--- naasi: 207
Methods and techniques: a point method of pasture analysis modified to determine proportion of blueberry plants in the Maritime Provinces affected by witches' brooms, 608; bioassay with chestnut blight fungus, Endothia parasitica, 672; formulae to calculate quantitatively the sporulation rate of an isolate of Helminthosporium maydis under a given set of environmental conditions, 730; a method to assess needle rust of balsam fir developed and used in Christmas tree stands 765; seed extraction with HCl and 3 hr drying at 66°C eradicated Corynebacterium michiganense in infected seeds, 974; tree girdling or felling to reduce oak bark beetle infection and fungus mat production in oaks, 1055
---: equipment: a device for monitoring the orifice direction of a Hirst spore trap, 244; automatically controlled movable shelters were designed to study effect of dew and rain on development of peanut leafspot epidemics, the moisture modification system can be used for other crops, 329; wind tunnel apparatus to measure effect of relative humidity and air speed on release of conidia of Helminthosporium maydis, 627; a modification of the Kramer-Collins spore sampler, 937
---: inoculation: efficient method for screening pear seedlings in greenhouse for resistance to fire blight, 20; inoculation of Trichoderma viride spores via pruning shears for biological control of Stereum purpureum on plum tree wounds, 25; a new pressure injector for forcing liquid fungicides and such into trees, is easily installed, light weight, and cheap, 338; a new trunk injection technique for systemic chemicals, 513; an improved technique for evaluating resistance of rice varieties to Xanthomonas oryzae, 537; a quantitative inoculation technique for screening sugarcane varieties for resistance to leaf scald, 582; inoculation, for evaluation of resistance in citrus seedlings to Phytophthora parasitica, 669; for mechanical transmission of citrus exocortis virus, 886; a method of cross-indexing pathogen isolates and host cultivars for virulence-susceptibility reactions, 957; spray inoculation compared with direct injection and forced-air chamber for mass inoculation of pine seedlings with fusiform rust, 1013;
---: for nematodes: simple extraction and use of hydrogen peroxide for estimating nematode populations in banana roots, 678; five methods of preparing inoculum of Meloidogyne spp. compared, 1025; a 4-minute lactophenol fixation method for nematodes, 1045
Mexico: 333
Michigan: 187; 230; 445
Microbial infection: of cotton seeds, vulnerability of seed to infection does not parallel that of the fiber at all times during preharvest weathering, 994
Milesia sp.: 765
Millet (Japanese, see Japanese millet)
Millet (pearl millet): Puccinia substriata var. indica found in widely separated areas on pearl millet in the Southeast in 1972, 262
Minnesota: 99; 353; 647; 890; 912
Mississippi: 59; 341
Mistletoe: dwarf, on lodgepole pine in Montana, new record, 298; detected on jack pine with aerial photographs, 951
---: European, Viscum album, additional hosts found in California, 904
Moisture: free moisture found to be necessary to obtain control of ball moss on shade trees with metallic fungicides, 81; its role in the sporulation of a race T isolate of Helminthosporium maydis, 304
Monilinia fructicola: 62; 457; 814
Monochaetia sp.: 522
--- unicornis: 315
Monoculture: of soybeans for 10 years in Minnesota, roots, rhizosphere, plant residues, and soil yielded five species of Fusarium, 912
Montana: 298; 701
Morphological description: conidial, of new race of Helminthosporium carbonum, 822
Morphology: colony morphology of Xanthomonas oryzae revealed two distinct types, a yellow and a white type, 955
Moth bean (Phaseolus aconitifolius): host of yellow mosaic disease, 800
Mountain lover (Pachistima myrsinites): sporophores of Fomes annosus found attached to stems, 407
Mountain mahogany (Cercocarpus sp.): Fomes annosus found on, 407
Mowing: effect of nitrogen fertilization and

mowing on Helminthosporium leaf spot of bluegrass leaves, 544
Mung bean (Phaseolus aureus): mung bean yellow mosaic virus, its characteristics, incidence and host range in India, 463; is susceptible to yellow mosaic disease that affects urd, moth and soybeans in West Pakistan, 800
Mustard (Brassica juncea, see Cruciferae)
Mycocentrospora sp.: 807
Mycological description: of Corynespora cassiicola on leaves of paper-mulberry, 296; of Taphrina deformans, cause of apricot leaf curl, 361; of Septoria halophila from Hordeum leporinum, 367; of Corynespora cassiicola on tobacco, 375; of Pithomyces chartarum from rice, 436; of Colletotrichum dematium from strawberry, 445; of Mycocentrospora sp., pathogen of Euonymus, 807; of Coryneum beijerinckii on peach in Iraq, 855
Mycoplasma: greening disease of citrus, caused by a mycoplasmalike organism (MLO), young lime trees injected with B. P. -101 showed excellent recovery and produced normal foliage and fruits, 160; occurrence of a spiroplasma in infected corn in Mexico, that closely resembles organisms previously described in stunt-infected corn, 333; in American elm that was infected with elm phloem necrosis, 341; control of pear decline with a tetracycline, 634; mycoplasmalike organisms in phloem of palms in Florida affected by lethal yellowing, 861
Mycosphaerella citri: 294; 691
--- ligulicola: 624
Myzus persicae: 552; 841

Nandina domestica: cucumber mosaic virus causes leaf narrowing and brilliant wine-red coloring of new leaves, but is apparently not seed-transmitted, 917
Near-ultraviolet radiation: a substance "P310" appears to be closely related to near-UV and is also closely associated with sporulation in Botrytis cinerea, 760
Nebraska: 111; 291; 681; 873
Needlecast: brown spot needlecast in Scotch pine Christmas tree plantations in north-central States, caused by Scirrhia acicola, 55; of Scotch pine, caused by Lophodermium pinastri, 515; Swiss needlecast of Douglas-fir caused by Phaeocryptopus gäumanni reported for first time from Pennsylvania is the southern-most known distribution in Eastern North America, 984
Nematicides (see also Antibiotics, Chemicals, and Fungicides): several tested together: systemic nematicides used as seed treatment alone were ineffective in tests at two locations for control of root-knot nematode in cotton, 129; carbon disulfide, methyl bromide and 1, 3-D tested for preplant fumigation for grape rootstocks, 416; oxamyl and BAY 68138 controlled foliar nematodes of Rieger begonia, 603; oxamyl and aldicarb were effective for control of Aphelenchoides fragariae of begonia, 1015
---: aldicarb, Heterodera schachtii-infested soil treated with aldicarb resulted in greater sugarbeet yields than other nematicide treatments, 942
---: BAY 25141, as chemical bare root dip gave complete control of root-knot-infected peach trees, 489
---: 1, 3-D: preplant fumigation did not control nematodes and oak root fungus in California test in vineyard for replants, 115; deep placement of 1, 3-D and methyl bromide gave excellent initial control of root-knot and other nematodes in grape vineyard replants, in California, 619
---: DD: DD and methyl bromide controlled root-knot nematodes in greenhouse-grown cucumber and increased yields of cucumbers, 1032; DD and DD-MENCS controlled root-knot nematodes on tomato and increased yield, 1033
---: ethoprop, used in combination with disulfoton and trap-crop to control cowpea mosaic virus, 684
---: methyl bromide: preplant fumigation at 400 lb/A gave good control of nematodes and oak root fungus in vineyard replants in California, 115; deep placement in California vineyards controlled root-knot nematodes, 619; methyl bromide and DD were more effective than SMDC in control of root-knot nematode in greenhouse-grown cucumber, 1032
---: oxamyl: as soil drench controlled Pratylenchus penetrans in birdsfoot trefoil, 237; oxamyl and BAY 68138 were effective for control of foliar nematodes in Rieger begonia, 603
---: thionazin: controlled Meloidogyne incognita-infected peach trees with bare root dips, 489; thionazin, oxamyl, and BAY 68138 were not effective as bare root dips for control of Pratylenchus penetrans in raspberry root cuttings, 492
Nematodes: soil fumigation for control of nematodes and Armillaria mellea in vineyard replants with methyl bromide, 115; loss of cotton yield from nematodes in the U. S. for a 7-year period has increased, 325; Aphelenchoides cibolensis and A. composticola, mycophagous nematodes, apparently reduced vigor or growth of Armillaria mellea root rot of

Nematodes (continued): Pinus ponderosa but neither killed the fungus, 355; association of nematodes and fungi in the root systems of bananas in state of decline in Lebanon, 615; incubation of banana roots in hydrogen peroxide for extraction of nematodes, 678; method described for lactophenol fixation of nematodes in 4 minutes, 1045
---: cyst: Heterodera latipons found on winter oat crop in Israel, 53; Heterodera mothi found for first time in the United States in Georgia on roots of yellow nutsedge, 946; Heterodera schachtii-infested soil treated with aldicarb produced greater sugarbeet yields than other treatments, 942
---: foliar: chemical control of Aphelenchoides fragariae on Rieger begonia, 603; A. fragariae controlled on Rieger begonia with oxamyl and aldicarb, 1015
---: grass root-gall nematode, Ditylenchus radicicola in western Oregon, its distribution and host range, 1021
---: Hoplolaimus seinhorsti, parasitic on cowpea in Nigeria, 798
---: lesion: Pratylenchus penetrans, new hosts of this nematode in Honduras and Panama plus a list of recorded hosts, 47; is new nematode pest in Trinidad, 260; controlled in raspberry root cuttings by hot water treatments, 492; reproduction in roots of seven cover crop species in greenhouse experiments, 728
---: needle: Longidorus cohni is winter pest of oats in Israel, 53
---: plant-parasitic: nutsedges that have increased in population in cotton fields with the use of herbicides are hosts to two nematode genera, 402; preplant fumigation for grape root stocks against plant-parasitic nematodes, 416; soil amendments of mycelial residues, cellulosic wastes and chitin reduced populations of Pratylenchus penetrans and Tylenchorhynchus dubius, 438; plant-parasitic nematodes associated with Caladium in Florida, 558
---: reniform: tomato resistance and susceptibility to Rotylenchulus reniformis, 169; distribution, occurrence, and seasonal occurrence in Georgia, 399
---: root-knot: root-knot and other nematodes controlled in California vineyards with deep placement of 1, 3-D and methyl bromide, 619; collecting inocula of Meloidogyne spp. by several methods, 1025; Meloidogyne hapla on potatoes in West Pakistan, a first report of the occurrence here, 505; M. hapla, screening alfalfa seedlings for resistance is best done at optimum inoculum levels of 600-800 larvae per plant, 657; systemic nematicides as seed treatment alone on cotton, did not give control of M. incognita acrita in Texas trials, 129; M. incognita races found in Louisiana that show variability in pathogenicity to sweetpotato, 199; sweetpotato bedding roots infected with M. incognita served to transmit the nematodes to the field via sprouts, 371; M. incognita controlled in root-knot infected peach trees with bare root dips of thionazin and BAY 25141, 489; all of U. S. D. A. plant introductions and several Japanese hybrids are susceptible to M. incognita, 606; M. incognita is important pathogen of tomato in Haiti, 921; methyl bromide and DD gave control of M. incognita in greenhouse-grown cucumber in Greece, 1032; M. incognita controlled on tomato with DD and DD-MENCS, in Greece, 1033; several tomato varieties field-tested for their reaction to M. javanica bearing the Mi gene for resistance showed no galling, two were very susceptible, 141; M. naasi was not pathogenic to 10 soybean cultivars screened for susceptibility, 207
---: sting, populations of Belonolaimus longicaudatus from North Carolina and Georgia have wide host ranges, 750
---: stylet, several together, isolated from turfgrass sod production fields in Pennsylvania, 891
Net blotch: selecting resistance to net blotch in Lolium-Festuca hybrids, 563
New diseases: glume blotch of rice, caused by Pithomyces chartarum, 436; basal gall of sunflowers incited by Plasmopara halstedii, 647; Mycocentrospora leaf spot of Euonymus in Ohio, 807
New distribution: downy mildew and rust found on St. Augustinegrass in Louisiana, 16; of comandra blister rust in western Kentucky, 17; of Sirococcus tip dieback of Pinus spp. in California forest nurseries, 69; bacterial wilt of carnation found in New Zealand, 136; Itersonilia petal blight of chrysanthemum found in Florida, 181; Sirococcus strobilinus is reported for the first time in the Lake States and the first time on red pine, 246; Pratylenchus penetrans found in Trinidad, West Indies for first time, 260; Puccinia substriata var. indica confirmed in the United States on pearl millet in several locations in the Southeast, 262; apparently first report of Cronartium comandrae on mugo pine in North America, 283; Goss's bacterial wilt of corn has spread to one county in Kansas, 291; of dwarf mistletoe on

lodgepole pine in north-central Montana, 298; first report of Scleroderris canker in New England, in Vermont, 338; apricot leaf curl, caused by Taphrina deformans, found in India, 361; Septoria halophila found to be a pathogen of Hordeum leporinum in Western Australia, first report in literature since it was first described in Argentina, 367; Meloidogyne hapla found in galls on several potato varieties in West Pakistan for the first time, 505; seedborne Pseudomonas glycinea in soybean seed in India, 531; bacterial leaf spot of zinnia caused by Xanthomonas nigromaculans f. sp. zinniae in Ohio, 555; Dutch elm disease moves into Montana, 701; Hoplolaimus seinhorsti recorded for first time from African continent and also as a parasite of cowpea, 798; peanut rust found in Australia for first time, 803; bacterial wilt of cassava found in Nigeria, 824; first record of outbreak of Coryneum blight on stone fruits in Iraq, 855; Heterodera mothi reported from Georgia, on nutsedge, 946

New hosts: of Pratylenchus coffeae from Central American banana-producing areas, including a list of previously recorded hosts, 47; colonial bentgrass cultivar Exeter is susceptible to Drechslera gigantea, 84; Fusarium lateritium f. pini on eastern white pine in Virginia, 182; Sirococcus strobilinus on red pine in the Lake States, 246; tomato seedlings infected by Pythium arrhenomanes in south Florida, thought to be first report, 260; Cronartium comandrae found on mugo pine in Alberta, Canada, 283; of European mistletoe (Viscum album) found in California, 904

New Jersey: 487

New York: 457; 468; 688; 817

New Zealand: 136

Nicotiana tabacum (see Tobacco)

Nigeria: 375; 798; 824

North Carolina: 325; 387; 551; 750; 845; 1020

North Dakota: 449; 647

Northern jointvetch (Aeschynomene virginica): newly discovered anthracnose disease of this rice weed was found to be endemic in 31 rice-growing areas in Arkansas, 695

Nuclear response: nuclei of Diplodia-infected corn increased in size and mass in response to inoculation and infection, 1043

Nutsedge (Cyperus esculentus and C. rotundus): as hosts of plant-parasitic nematodes in Georgia cotton fields, 402; C. esculentus is host of Heterodera mothi, 946

Oak: black and blackjack, association of viruslike particles with a ringspot disease, 1029

---: wilt: recovery of compatibility types of gray and albino strains of Ceratocystis fagacearum from paired culture inoculations of Quercus coccinea, 254; new isolation in North Carolina shows spread north and east of previously known infections, 325; density of oak bark beetles was lowest where diseased trees were felled; the incidence of oak wilt declined when density of beetle attacks dropped, 1055

Oak root fungus (Armillaria mellea): preplant fumigation with methyl bromide in vineyard replants at 400 lb/A under polyethylene cover appears promising treatment for control of A. mellea, 115

Oats: two nematode species are causing economic damage to winter oats in Israel, 53; stem rust, races 31, 61 and 32 were most important in the USA during 1972, 754

Ohia (Metrosideros collina ssp. polymorpha): Phytophthora cinnamomi, cause of root rot and death of ohia in Hawaii, is correlated with high rainfall and poor drainage, but not with ohia decline, 383

Ohio: 156; 515; 555; 807

Oidium sp.: 475

Onion: 'Buccaneer' had lowest incidence of Botrytis brown stain of 11 yellow onion cultivars tested at harvest and after 4 months' storage, 210

Ophiobolus graminis: 187

Oregon: 94; 716; 1021

Orobanche ramosa (branched broomrape): caused a definite yield reduction in parasitized tomato plants in California, 926

Ovatus crataegarius: first report as a vector of pea seedborne mosaic virus, 552

Overwintering: pea seedborne mosaic virus was shown to overwinter effectively in seedlings of hairy vetch, 349

Oxamyl (see Nematicides)

Ozone: ozone injury to tobacco plants in South Dakota, 106; potato varieties showed differential resistance to ozone injury, 542; 'Golden Jubilee' commercial sweet corn displayed significant response to ozone exposure in plant height, length of internodes and fresh and dry weight, 719; poinsettia cultivars resistant to ozone injury, 774; response of nine forest tree species to chronic ozone fumigation, physiological measurements as well as foliar symptoms should be considered, 914

"P310" close association of this substance

with both near-ultraviolet radiation and sporulation of Botrytis cinerea, 760
Palm: lethal yellowing, caused by mycoplasmalike organisms found in fan palm and Christmas palm in Florida, 861
Panama: 47
Papaya (Carica papaya): postharvest rots were reduced in green papaya fruit when treated with benzyl isothiocyanate as a dip or as a vapor, 86; a stem and fruit disease in Barbados caused by Corynespora cassiicola, 404; leafhopper Empoasca stevensi found as new vector of bunchy top disease of papaya, 412
Paper-mulberry (see Broussonetia papyrifera)
Paratylenchus sp.: 891
Patch disease: of two bentgrasses, caused by Drechslera gigantea, in Rhode Island, 84
Pathogenicity: conidial morphology and pathogenicity of Alternaria alternata were influenced by culture medium composition, but the two characteristics were not related, 39; of Fusarium oxysporum to variegated pyracantha, 389; of Rhizoctonia solani and Pythium myriotylum from rotted pods to peanut seedlings, 578; of Verticillium dahliae to potato, 735
Pea: seed rot and damping-off, caused by Pythium ultimum, synergistic effect of chloroneb and thiram in seedling protection 3; several copper-containing compounds added to greenhouse soil suppressed Aphanomyces root rot, 876
---: virus: pea seedborne mosaic virus was transmitted to pea seed both by the female parent and by pollen, 248; PSbMV was naturally transmitted from infected vetch plants to susceptible pea cultivar Dark Skin Perfection planted nearby in two successive growing seasons in Wisconsin, 349; natural infection of pea by lettuce mosaic virus found in New York, no resistance found in 125 plant introductions, 688
Peach: fungicide-amended wound dressings to control Cytospora cincta, 28; silver leaf disease, caused by Stereum purpureum, was found to be widespread in California peach orchard in 1971, 192; chemical bare-root dips in BAY 25141 and thionazin completely controlled root-knot-infected trees, 489; fungicidal effectiveness of hypochlorous acid at 0 to 25°C on Monilinia fructicola conidia, 814
Peanut (Arachis hypogaea): movable shelters constructed to study effect of dew and rain on development of leafspot of peanut worked satisfactorily for three seasons, 329; Rhizoctonia solani and Pythium myriotylum from rotted pods are pathogenic to peanut seedlings, 578; peanut rust found in the Northern Territory of Australia in April 1973, its first appearance on the continent, 803
---: Cylindrocladium black rot: its status in Georgia since its discovery in 1965, 90; over 40 infestations of Cylindrocladium black rot of peanuts were confirmed in North Carolina in 1972, 387; rotational crops corn and tobacco are susceptible to C. crotalariae, corn is resistant, 1035
Pear: fire blight, method developed to screen large numbers of pear seedlings for fire blight resistance, 20; pear vein yellows virus is systemic, is widespread in the Yakima Valley and is apparently spread through horticultural propagation, 483; pear decline, probably caused by a mycoplasmalike organism, was markedly reduced with oxytetracycline HCl transfusions, 634
Pearl millet (see Millet)
Pecan: control of pecan scab with triphenyltin hydroxide and benomyl, 902
Pelargonium (geranium): root and stem rot of cuttings by Rhizoctonia and Fusarium, 177; a disease with "diverse symptoms" previously thought to be caused by Verticillium albo- atrum was found to involve two bacterial isolates and a Fusarium isolate, 284; Pelargonium rust found in Minnesota in March 1973, 890; Pseudomonas is seed-transmitted in Pelargonium with "diverse symptoms" and is considered a contributory factor in the disease, 900
Pennsylvania: 104; 411; 563; 891; 984
Pepper: bacterial spot, found for first time on peppers in California, 258; further pepper virus identification in Florida, 991
Pestalotia phoenicis: 856
Pestalotiopsis palmarum: 856
pH: high aqueous-extract pH (an indicator of exposure to humid conditions) was not characteristic of Aspergillus flavus-infected cotton fibers, indicating A. flavus boll rot tends to occur in drier areas than boll rots caused by most other fungi, 664; relation of aqueous extract of cotton fiber, to location of cotton production, and resulting presence of certain fungi, 756; cotton seed infection can occur without increases in fiber pH and high fiber pH is not always accompanied by seed infection, 994
Phaeocryptopus gäumanni: 984
Phaseolus (see also Bean): germplasm tolerant to Nebraska strains of the common blight bacterium of bean found in several lines and cultivars of several Phaseolus species, 111

--- aconitifolius (see Moth bean or Bean)
--- aureus (see Mung bean or Bean)
--- mungo (see Urd bean or Bean)
Phialophora gregata (Cephalosporium gregatum): 269; 660; 853; 1063
Phloem: mycoplasmalike organisms found in phloem of two palms other than coconut in Florida, 861
Phloem necrosis: of elm, tetracycline antibiotic suppressed symptoms in American and cedar elms, 341
Phoma sp.: 856
--- betae: 204
Phomopsis cocoes: 856
--- juniperovora: 59; 315
Phyllosticta antirrhini: 592
Phymatotrichum omnivorum: 318; 403
Physalospora obtusa: 158; 676
--- tucumanensis (Colletotrichum falcatum): 220
Physiological diseases: pole blight of western white pine in Idaho, 948
Physoderma maydis: 630
Phytophthora cinnamomi: 383; 1005
--- citrophthora: 391
--- cryptogea: 190
--- infestans: 612
--- palmivora: 86; 529; 669
--- parasitica var. nicotianae: 832; 884
--- phaseoli: 724
Phytotoxicity: of dodine and azinphosmethyl to cherry fruit, 428
Picea rubens (red spruce): first report of Rhizina undulata on large trees in North America, on 80-year-old spruce in Vermont, 394
Pinus spp.: Sirococcus dieback in California forest nurseries, 69; root diseases in pine nurseries in the Australian Capital Territory, 1058
--- banksiana (jack pine): dwarf mistletoe can readily be detected on conventional vertical aerial photographs, 951
--- contorta (lodgepole pine): a new distribution record for dwarf mistletoe in Montana from the Bearpaw Mountains, 298
--- elliottii var. elliottii (slash pine): soil drenches of benomyl, oxycarboxin and CW-524 in greenhouse tests eradicated fusiform rust fungus, 776; spray inoculation with Cronartium fusiforme basidiospores for mass inoculation of seedlings for resistance tests, 1013
--- monticola (western white pine): pole blight found for first time in a plantation, in northern Idaho, 948
--- mugo (mugo or Swiss mountain pine): found as a new host for comandra blister rust, in Alberta, Canada, 283
--- ponderosa (ponderosa pine): two mycophagous nematodes retarded root rot development by Armillaria mellea on pine seedlings, 355
--- resinosa (red pine): Sirococcus strobilinus is cause of shoot blight in Lake States, a new host, 246; four fungicides were effective in control of Lophodermium pinastri, cause of needlecast disease in nursery seedlings, 263; Scleroderris canker found in plantation in Vermont, 338
--- strobus (eastern white pine): found to be a new host for Fusarium lateritium f. pini, pathogen isolated from a well developed canker, thought to be the first reported occurrence of the fungus on eastern white pine, 182
--- sylvestris (Scotch or Scots pine): brown spot needlecast, caused by Scirrhia acicola, is causing major defoliation in Scotch pine Christmas tree plantations in six north-central States, 55; 12-year-old pines found infected by Scleroderris lagerbergii in Vermont, 338; a mid-August and mid-September maneb spray recommended for practical control of Lophodermium needlecast of Scotch pine in Ohio, 515
--- taeda (loblolly pine): comandra blister rust found in 6-year-old plantation in Kentucky for the first time, 17
Piperazine fungicide (see Fungicides, Cela W 524)
Pisum sativum (see Pea)
Pithomyces chartarum: 436
Planetree (Platanus acerifolia): a tree wound dressing with benomyl added prevented spread of the Ceratocystis causing canker stain disease, 28
Planting date: of soybeans in India had significant effect on seed quality, 770
Planting rate: the effect of rate of planting on incidence of brown stem rot in soybean, 269
Plasmopara halstedii: 647
Plum: control of Stereum purpureum (cause of silver leaf disease) in pruning wounds by applying Trichoderma viride with pruning shears, 25
Poa pratensis (Kentucky bluegrass, see also Grasses): stripe rust of Kentucky bluegrass controlled with carboxin and oxycarboxin, 217
Poinsettia (Euphorbia pulcherrima): five commonly grown cultivars are relatively resistant to foliar ozone injury normally encountered in the Northeast, 774
Poland: 31
Pole blight: of western white pine, a disease resulting from a complex interaction of physiologically related factors, 948
Poor drainage: as a possible factor in increased infection of yellow birch

seedlings to Diaporthe cankering, 230
Poria weirii: 346
Postharvest decay: of blueberries from New Jersey in marketing channels, 487
---: of carrots: fungicide dips of benomyl and DCNA significantly reduced losses from rots, 697; as affected by bacterial contamination and refrigeration temperatures, 896
---: of citrus fruits, control with benomyl, thiabendazole and thiophanate methyl, 899
---: of mango, control of postharvest decay of fruit in South Africa with benomyl, 173
---: of papaya, inhibition of fungal pathogens of papaya fruit with benzyl isothiocyanate (BITC), 86
---: of raspberry, control of Botrytis cinerea on fruits with acetaldehyde vapor, 795
---: of snap beans, hot water and DCNA dips controlled Pythium and Sclerotinia rots, 234
---: of stone fruits, brown rot (Monilinia fructicola) control with heated fungicidal dips, 62
---: of strawberry, berries held in CO_2-enriched atmospheres had much less softening and decay than fruit held in air, 44
Potato: root knot on potato in West Pakistan, 505; in variety tests in Delaware, 'Superior' was most resistant to ozone injury and Haig the most susceptible, tests suggest the value of testing breeding lines for resistance to air pollution as a standard procedure, 542; isolate of Verticillium dahliae is pathogenic to commercially grown potatoes in Australia, 735; Rhizoctonia and blackleg, fungicidal seed and soil treatments reduced symptoms, 803; Verticillium species caused no severe damage to potatoes from seed sources from the northern Northeast grown in Rhode Island, whereas four varieties developed in North Dakota were severely damaged by V. albo-atrum, 928
Powdery mildew: of wheat, in Michigan, 187; of Cassia occidentalis (a local lesion host for Cymbidium mosaic virus), benomyl gave complete control of the powdery mildew, 475
Pratylenchus sp.: 891
--- coffeae: 47
--- penetrans: 237; 260; 438; 492; 728
--- vulnus: 416
Preharvest infection: of U. S. cotton fiber in relation to location of growth and to aqueous-extract pH of fiber, 756
Prince Edward Island: 237
Pritchardia pacifica (fan palm): found infected with lethal yellowing in Florida, 861
Prunus (see also common names): benomyl plus DCNA in combination with hot water in 3-minute dips was more effective than fungicidal dips with cold water for control of brown rot of prune fruit, 62; Prunus ringspot virus transmitted by pollen in prune and sour cherry orchards in Oregon, 241; Coryneum blight of stone fruits in Iraq, 855
Pseudomonas sp.: 284; 909
--- cichorii: 504
--- glycinea: 156; 531
--- tomato: 258
Pseudopeziza medicaginis: 468
Pseudopityophthorus sp.: 1055
Pseudotsuga menziesii (Douglas-fir): dwarf mistletoe infection sites west of the Cascade Range in Oregon are raised from two to nine, 716; Swiss needlecast on Douglas-fir in Pennsylvania, 984
Puccinia spp.: 793
--- antirrhini: 709
--- arachidis: 803
--- coronata: 215
--- epilobii: 765
--- graminis: 215; 595
--- --- f. sp. avenae: 754
--- --- f. sp. tritici: 8; wheat stem rust epidemic potential in 1972, 434; 567; 880
--- helianthi: 524
--- pelargonii-zonalis: 890
--- recondita: 567
--- --- f. sp. tritici: 8; 187; 449; 567; 572; 653
--- stenotaphri: 16
--- striiformis: 8; 215; 217
--- substriata var. indica: 262
Pucciniastrum goeppertianum: 608; 765
Puffballs (Lycoperdon gemmatum): BAS 3191 F, Dyrene, and carboxin at low concentrations inhibited growth of fairy ring puffball fungus in bentgrass turf, 711
Pulse crops (grain legumes): symptomatology, transmission, host range, incidence, losses and control of four viruses affecting urd and mung beans in India, 463
Pyracantha (variegated): Fusarium oxysporum caused wilting of variegated Pyracantha but not of the common form; benomyl as a soil drench provided control, 389
Pythium spp.: 403; 1058
--- aphanidermatum: 638; 1066
--- arrhenomanes: 260
--- butleri: 234
--- irregulare: 848
--- myriotylum: 260; 578
--- sylvaticum: 848
--- ultimum: 3; 848; 1066

Queensland (see also Australia): 1040

Queensland maple (Flindersia brayleyana): Rhizoctonia disease of seedlings was controlled by seed coating of PCNB, 907
Quercus coccinea (See Oak)

Races: seven physiologic races of Colletotrichum lindemuthianum identified on Phaseolus vulgaris in Minas Gerais, Brazil, 870
---: Helminthosporium: new Helminthosporium described from the Corn Belt, 780; new race, designated race 3, of H. carbonum, 822
---: --- maydis: races O and T: further studies on the past occurrence and geographic distribution of isolates of race T, 18; at least 4 hours of high humidity required for significant lesion production on corn seedlings, 76; influence of climatic factors on colonization of a susceptible corn hybrid by race T, 145; studies show that previous isolates of race T in the U. S. possessed the ability to attack all aboveground parts of corn hybrids with male-sterile cytoplasm, 253; spore sampling of race T, 298; influence of some climatic factors on sporulation of race T, 304; a corn seed lot infected with race T showed improved seedling stand over the control when treated with carboxin + thiram, 344; racial composition of Helminthosporium maydis on corn hybrids in normal cytoplasm in Pennsylvania in 1972, 411; races O and T and race 2 of H. carbonum found capable of inducing infection on pop corn, johnsongrass, barley, rye, and pearl millet, 507; electrolyte and peroxidase leakage used as indicator of susceptibility of maize inbreds to H. maydis, 518; cross-protection by race O against race T, 971; incidence of race O and T on corn with normal and Texas male-sterile cytoplasms at Gainesville, 1971-1972, 981
---: of Phytophthora parasitica var. nicotianae race 1, in Georgia, 832
---: of Puccinia graminis f. sp. avenae in the USA in 1972, 754
---: of P. graminis f. sp. tritici in 1972, 880
Radiation (see Near-ultraviolet radiation)
Ramulispora sorghi: 681
Raspberry: recovery of tomato ringspot virus from seedlings indicated seed transmission to seed via maternal tissue, 431; control of Pratylenchus penetrans in root cuttings with hot water dip at 118-119°F for 15 minutes, 492; postharvest decay control with acetaldehyde vapor, 795
Ratoon stunting disease: of sugarcane, diagnosis based on presence of a specific bacterium, 987
Relative humidity: high-low RH with increased air speed increased the spore release of Helminthosporium maydis conidia in wind tunnel experiment, 627
Reproduction: and host range of Belonolaimus longicaudatus, 750
Resistance: of tomato to the reniform nematode, 169; in Beta vulgaris to Phoma storage rot, selection for resistance should be done at 20-25°C, 204; sources of resistance to a soilborne fungal disease complex of sunflowers, 318; of Medicago spp. (alfalfa) to leafspot fungus, Pseudopeziza medicaginis, 468; in American and Indian barley cultivars to Helminthosporium stripe, 495; to rust and Verticillium wilt in sunflower varieties, 524; Lolium-Festuca hybrid clones 57-3 and 68-1 showed resistance to Helminthosporium dictyoides, 563; monogenic and multigenic resistance to Helminthosporium turcicum in corn, 586; of Poa pratensis introductions to Puccinia graminis, 595; to Phytophthora parasitica in citrus seedlings, 669; to Phytophthora in kenaf and related species, 884; in 'Apalachee' alfalfa to Stemphylium leafspot in field and greenhouse tests in North Carolina, 946; in Chinese elm hybrids to Dutch elm disease, 997; some resistance in corn cultivars to southern corn leaf blight resulted in smaller lesions and fewer conidia than on susceptible cultivar, 1047
Rhizina undulata: 167; 394
Rhizoctonia solani: 177; 222; 403; 438; 453; 549; 578; 601; 615; 642; 803; 907
Rhizopus stolonifer: 86; 756
Rhode Island: 84; 928
Rhopalosiphum padi: 552
Rhynchosporium oryzae: 715
Rice: a new glume blotch of rice caused by Pithomyces chartarum, found for first time in India, 436; leaf scald, caused by Rhynchosporium oryzae, in Louisiana, 715
---: bacterial leaf spot: lesion expansion of Xanthomonas oryzae was faster in susceptible than resistant varieties of rice and pathogen lives and multiplies mostly in living parts of the rice plant tissue, 227; a new clipping technique for evaluating resistance of rice varieties to X. oryzae, 537; different rates of movement and multiplication of bacterium occur in susceptible and resistant hosts, 784
Robinia pseudo-acacia (black locust): a virus resembling tomato spotted wilt virus found in black locust in Arkansas, 153
Root diseases: of pine seedlings in nurseries in Australia, 1058
Roselle (Hibiscus sabdariffa): several

introductions and breeding lines showed resistance to Phytophthora parasitica nicotianae, 884
Rot: Alternaria rot, of blueberries, 487
---: boll rot, of cotton, yield loss from boll rots has increased in the U. S. from 1965-1971, 325; Aspergillus flavus boll rot tends to occur in drier areas than most other boll rots, 664
---: brown rot, of grapefruit following sprinkler irrigation with Phytophthora-infested water, 391; of stone fruits, CW-524 proved to be a very promising new fungicide for the control of brown rot, caused by Monilinia fructicola, 457; of peach, effect of chlorinating hydrocooling water on Monilinia fructicola conidia and brown rot infection, 814
---: brown stem rot of soybean: planting rate affects the incidence of brown stem rot, 269; incidence in Iowa in 1972, 660; incidence development, and yield effects of Cephalosporium gregatum on soybeans, 853; a collection of isolates of Phialophora gregata varied in cultural type, but less in virulence or growth rate, 1063
---: charcoal rot of soybeans: causal organism, Macrophomina phaseolina, remained pathogenic after 9 months' storage in liquid nitrogen, 478; decreased with growth regulators, 789
—-: collar rot and wilt of kenaf and roselle caused by Phytophthora, 884
---: corm: of Colocasia esculenta (taro), by Fusarium oxysporum, 701; of gladiolus, caused by Fusarium, method developed for cross-indexing pathogen isolates and host cultivars for virulence-susceptibility reactions, 957
---: Cylindrocladium black rot of peanut: is now confirmed in eight counties in Georgia, 90; its incidence in North Carolina in 1972, 387; susceptibility of peanut rotational crops cotton and tobacco to C. crotalariae, 1035
---: foot rot: of wheat, caused by Fusarium spp., 187; foot and root, of citrus, by Phytophthora parasitica, 669
---: fruit: of mango, heated benomyl gave control, 173
---: ---: of tomato: Rhizoctonia solani, Sclerotium rolfsii and Erwinia sp. accounted for 85% of the organisms isolated from diseased tomatoes in Mississippi, 453; Pythium fruit rot of tomatoes in Florida, 638; caused by Pythium ultimum and P. aphanidermatum, in California, 1066
---: red rot of sugarcane, treatment of canes at 54°C for 8 hr killed red rot fungus and inactivated the causal agents of the grassy shoot and ratoon stunting disease of sugarcane, 220
---: red stalk rot of sorghum, caused severe damage to ratooned sorghum in Hawaii, 272
---: Rhizina root rot in conifer plantations in western Washington, 167
---: root: root and stem, of geranium cuttings, caused by Rhizoctonia and Fusarium, 177; of grand fir (Abies grandis), Poria weirii, Fomes nigrolimitatus, or Armillaria mellea occurred most frequently and with the most damage in Idaho, 346; of Pinus ponderosa by Armillaria mellea, two mycophagous nematodes retarded root rot development, 355; of ohia (Metrosideros collina), Phytophthora cinnamomi infection not correlated with ohia decline, 383; of pea, by Aphanomyces euteiches, suppressed with copper compounds, 876
---: seed, of pea controlled with chloroneb-thiram combination, 3
---: soft rot of carrots, caused by Erwinia carotovora, incidence was influenced by total bacterial populations in packing-house waters and postharvest holding temperatures, 896
---: stalk: of corn: caused by Diplodia maydis, cob parenchyma suggested as possibly an ideal tissue in which to study host-pathogen interactions of stalk rot organisms, 308; a first report of a positive response of corn nuclei to D. zeae by increase in size and mass, 1043
---: ---: of corn and sorghum, survival of Macrophomina phaseoli in stalk residue, 873
---: stem-end rot of papayas, reduced in mature-green fruits by treatment with benzyl isothiocyanate, 86
---: storage: rot of giant sequoia seedlings by Botrytis cinerea, controlled with benomyl, 67; of sugarbeet, of 1099 roots from the World Collection inoculated, 197 showed resistance to Phoma betae, 204
---: strawbreaker foot rot of winter wheat, causal pathogen Cercosporella herpotrichoides shows lack of benomyl tolerance, 935
Rotylenchulus reniformis: 169; 399
Rust: longevity of rust spores in liquid nitrogen, 793
---: blister: comandra blister rust caused by Cronartium comandrae found in western Kentucky on loblolly pine for the first time, 17; of mugo pine, first report on this host in North America, 283
---: of blueberry, production losses from rust fungus in the Maritime Provinces, 608
---: crown, Puccinia coronata found on perennial ryegrass in Hawaii, 215

Rust (continued): fusiform, Cronartium fusiforme eradicated from slash pine seedlings in greenhouse tests with three systemic fungicides, 776
---: leaf: of wheat: a study of the incubation period and rate of spread of leaf, stem, and stripe rusts in India indicate the importance of temperature as a limiting factor in the development and spread in northwestern India, 8; in Michigan during 1969-1972, 187; Puccinia recondita f. sp. tritici virulence in North Dakota in 1972, 449; leaf and stem, of wheat, results from seed treatment with oxycarboxin suggest that successful control by seed treatment with a systemic fungicide in the north-central States requires suppression of a rust epidemic for more than 80 days, 567; control of leaf rust of wheat in India with 4-n-butyl-1, 2, 4-triazole (BT) added in irrigation water or as a spray, 572; effect of dose, time of application, and variety of wheat on control of leaf rust by BT, 653
---: needle rust of balsam fir, use of a quantitative assessment method to assess needle rust in Maritime Provinces, 765
---: Puccinia antirrhini, growth in imbibed snapdragon seeds, 709
---: P. arachidis, on peanut, in Australia, the first appearance on the Australian continent, 803
---: P. helianthi, on sunflowers, varieties tested for resistance to rust and Verticillium wilt, 524
---: P. pelargonii-zonalis, on geranium, 890
---: P. stenotaphri, found on St. Augustinegrass in Louisiana for the first time, 16
---: P. striiformis: found on Kentucky bluegrass cultivars in Hawaii in 1972, a first report, 215; control on Kentucky bluegrass, with fungicides, 217
---: P. substriata var. indica, observed on pearl millet, in the Southeastern United States in 1972, 262
---: stem: of wheat, the epidemic potential in 1972 was averted by the nearly universal use of early maturing and resistant varieties and nearly complete eradication of barberries, 434; reaction of Poa pratensis introductions to the rust encompassed complete range of susceptibility, 595; of oat, important races in 1972 were races 31, 61, and 32, 754; of wheat, prevalence of races of Puccinia graminis f. sp. tritici in the USA during 1972, 880
Rye: reaction of rye addition lines in wheat to wheat streak mosaic virus, 6; found to be a good host of Pratylenchus penetrans, 728
Santolina spp.: Santolina blight, caused by Alternaria tenuissima, 1072
Scab: of apple, addition of summer superior oil to benomyl spray reduced percentage of scabbed leaves in South Australia, 560; tolerance of Venturia inaequalis to dodine in relation to history of dodine usage in apple orchards, 817
---: of pecan, caused by Fusicladium effusum, controlled with TPTH and benomyl, 902
Schizaphis graminum: 533
Scirrhia acicola: 55
Scleroderris lagerbergii: 338
Sclerophthora sp.: 16
Sclerospora sorghi: 42
Sclerotia: survival of sclerotia of Rhizoctonia solani in lake water, 601; reduction of sclerotia of Typhula incarnata by Trichoderma spp. to reduce gray snow mold of turfgrasses, 924
Sclerotinia sclerotiorum (= Whetzelinia sclerotiorum): 164; 234; 770; 932; 1070
Sclerotium rolfsii: 453; 770
Scolytus multistriatus: 511
Seed: increased recovery of Aspergillus flavus from soybean seed was correlated with decreased germination, 185; seed blight of snapdragon is caused by several seedborne fungi, 592; quality of soybean seed is affected by planting date, moisture conditions at harvest, and storage conditions, 770; uninfected seeds of Datura are brown, but Alternaria crassa-infected seeds were gray, 960; the process of microbial infection in cotton seeds in the field, 994
Seed: transmission: of southern bean mosaic virus in cowpea, 13; of pea seedborne mosaic virus in pea, 248; of tomato ringspot virus in raspberry, 431; of tomato ringspot virus in Gomphrena globosa, 433
Seed: treatment: of pea to control seed rot and damping-off by Pythium ultimum, 3; cotton treated with systemic nematicides alone did not give control, 129; carboxin + thiram applied to corn seed infected with Helminthosporium maydis race T significantly improved seedling stands, 344; benomyl and thiabendazole controlled Ascochyta blight of chickpea, 742; tomato seed extraction with HCl eradicated Corynebacterium michiganense, 974; none of several chemical seed treatments for 'Nugaines' winter wheat increased plant stand, yield, or test weight in eastern Washington and Oregon, 1006
Seedborne infection: Pseudomonas glycinea is seedborne on soybean in India, 531; seedborne fungi are cause of snapdragon seed blight, a serious problem, 592; of snapdragon seed with rust fungus, Puccinia

antirrhini, 709; fungi of soybean affect seed quality, 770; Alternaria crassa infects Datura and is seedborne, 960; bacterial leaf and flower spot of zinnia is apparently seedborne, 1020
Seedling: monocrotophos assists movement of fungicide carboxin in cotton seedlings, 642
Seedling disease: Botrytis storage rot of giant sequoia seedlings, 69; Sirococcus tip dieback of Pinus spp., 69; outbreak of Diaporthe cankering in 8-year-old yellow birch seedlings, infection found to be greatest on poorly drained sites, 230; control of needlecast disease of red pine nursery seedlings with fungicides, 263; cotton losses have increased in California over a 7-year period, 325; pathogenicity of Rhizoctonia solani and Pythium myriotylum from rotted peanut pods, 578; of corn to test changes in Helminthosporium turcicum following serial host passage, 590; three species of Pythium identified that cause seedling blight of cotton, 848; susceptibility of Queensland maple seedlings to Rhizoctonia disease was inversely correlated with the age of the seedlings, 907; severe, uniform response by monoembryonic 'Etrog' citron seedlings to citrus exocortis virus, 1010; root diseases of pine nursery seedlings, 1058
Segregation: factors for the two phases of early blight of tomato caused by Alternaria solani segregate independently of each other, 964
Septoria apiicola: 425
--- avenae f. sp. triticea: 99
--- halophila: 367
--- nodorum: 99; 187; 363
--- tritici: 99
Sequoiadendron giganteum (giant sequoia): benomyl as a prelifting field spray or a postlifting plant dip gave excellent control of Botrytis storage rot of seedlings, 67
Serial host passage: of Helminthosporium turcicum six times through monogenic resistant or susceptible corn seedlings did not change virulence of the pathogen, 590
Serology: of quail pea mosaic virus, 311
Sexual types: of Ceratocystis ulmi in Wyoming, 576
Sierra Leone: 509
Silver buttonwood (Conocarpus erectus var. sericeus): Cylindrocladium scoparium is causal agent of a new disease on this evergreen in Florida, 500
Silver leaf disease: of fruit trees, caused by Stereum purpureum, inoculation of Trichoderma viride spores via pruning shears controlled S. purpureum on plum tree wounds, 25; of peach, in California, first known occurrence in a commercial peach orchard in the State, 192
Sirococcus strobilinus: 69; 246
Smut: of sugarcane in Hawaii, caused by Ustilago scitaminea, 966
Snapdragon: seed blight, a serious problem caused by seedborne fungi, principally Alternaria alternata, 592; Puccinia antirrhini found seedborne in seedlot; seeds internally infected are suggested as a possible means of undetected disease being introduced into rust-free areas, 709
Snow mold: viability of sclerotia of Typhula incarnata drastically reduced after exposure to Trichoderma spp., 924
Soil: Xanthomonas alfalfae, bacterial leaf spot pathogen, found to overwinter in soil in Kansas, 52; incidence of Verticillium species in soils, vines and tubers of Rhode Island-grown potatoes, 928; excessive soil moisture associated with incidence of Santolina blight, 1072
---: amendments, industrial mycelial residues, wood fiber wastes and chitin added to soil reduced populations of plant parasitic nematodes and Rhizoctonia, mycelial residues made Fusarium wilt of tomatoes worse, 438
---: treatment: for fungi: pressure-injected benomyl + Tween 20 protected elms against Dutch elm disease, 35; benomyl as a soil drench gave effective control of cutting decay of two foliage plants, Gynura and Hoya, 222; soil drench fungicides ineffective in control of Cylindrocladium scoparium on silver buttonwood cuttings, 500; ethoprop + disulfoton gave highest cowpea yields in trap-cropping experiment to control cowpea mosaic virus, 684; soil drenches of benomyl, oxycarboxin and CW-524 eradicated fusiform rust, 776; with copper compounds to suppress Aphanomyces root rot of peas, 876; with fumigants and fungicide drenches for control of Sclerotinia sclerotiorum in tobacco seedbeds, 932
---: ---: for nematodes: oxamyl controlled Pratylenchus penetrans on birdsfoot trefoil, 237; preplant fumigation with nematicides for grape rootstocks, 416; 1, 3-D and methyl bromide at high dosage and deep placement gave excellent initial control of root-knot nematodes in vineyards and for replants in California, 619; nematicidal treatments against Heterodera schachtii, 942; with several pesticides to test their control of foliar nematode on begonia, 1015; nematicides methyl bromide and DD controlled Meloidogyne incognita in greenhouse-

grown cucumbers, 1032; with nematicides for control of root-knot nematode on tomato, 1033
Soilborne diseases: effect of industrial mycelial residues, wood fiber wastes and chitin on Fusarium sp. and Rhizoctonia solani, 438
Solanum melongena (see Eggplant)
Sooty stripe: of sorghum, its occurrence in Nebraska may be the most northern record of its distribution, 681
Sorghum (Sorghum bicolor): a mosaic disease affecting several grasses, including corn and sorghum is reported from Israel for the first time, 125; Colletotrichum graminicola, caused severe stalk rot damage and lodging in Hawaii in 1971 and 1972, 272; three mechanically transmissible viruses found in sorghum and corn in South Dakota in 1972, 373; a severe foliar blight caused by Rhizoctonia solani, in Georgia, 549; sooty stripe, has been found in Nebraska, overwintering primarily as sclerotia, 681; Macrophomina phaseoli sclerotia survived in stalk residue for 18 months in Nebraska, 873
South Africa: 173; 1005
South Carolina: 917
South Dakota: 106; 373
Southern corn leaf blight (see Corn)
Southern pea (see Cowpea)
Soybean: recovery of Aspergillus flavus from soybean seeds was negligible at 20°C, but was recovered from 17 of 20 lots at 35°C, 185; 10 cultivars screened for susceptibility to the barley root-knot nematode, Meloidogyne naasi, were poor hosts, 207; about 45 days after seeding, Rotylenchulus reniformis builds to high levels, 399; seed quality as it is affected by planting date, storage conditions, and seedborne fungi, 770; five Fusarium species isolated from roots, rhizosphere, and plant residues of soybean and from soil in Minnesota, 912
---: bacterial blight: Pseudomonas glycinea associated with diseased plant parts overwintered on soil and on sod in Ohio, 156; seedborne P. glycinea and fungi affect soybean seed quality in India, 531
---: brown stem rot: incidence of brown stem rot increased as the planting space within the row was increased during 3 successive growing seasons, 269; caused by Cephalosporium gregatum, incidence in Iowa in 1972, 660; incidence, development and yield effects of C. gregatum on soybean in Illinois, 853; variation in Phialophora gregata, 1063
---: Macrophomina phaseolina: two strains isolated from soybeans and stored in liquid nitrogen retained their pathogenicity for 9 months, 478; growth regulator treatments reduced severity of charcoal rot of soybean, 789
---: virus: aerial detection of tobacco ringspot virus-infected soybean plants, 471; host of yellow mosaic disease in West Pakistan that is transmitted by whitefly, 800; economic injury thresholds of bean pod mottle virus and tobacco ringspot virus infection of soybean, 811; bean pod mottle virus, healthy plants yielded higher than BPMV-infected plants irrespective of various soil water regimes; water stress with infection further decreased yield, 1050
Spacing: wider row spacing and determinate plant growth in Great Northern beans reduced amount of white mold disease, caused by Whetzelinia sclerotiorum, 1070
Spain: 649
Spiroplasma: spiral-shaped microorganisms observed in corn stunt-infected plants in Mexico, 333
Spore age: of Septoria nodorum, and the effect on virulence, 363
Spore release: of Lophodermium pinastri, cause of needlecast of red pine, fewer fungicidal sprays are necessary for control if spray application is correlated with spore release, 263
Spore sampling: of Helminthosporium maydis race T in seven States in 1971, 298; a modification of the Kramer-Collins spore sampler, 937
Spore trap: a Hirst spore trap was modified to facilitate a continuous record of the orifice direction, 244
Sporulation: three formulae have been developed to predict the sporulation rate of Helminthosporium maydis lesions over a range of dew lengths and dew temperatures, 730; substances from Botrytis cinerea associated with sporulation and exposure to near-UV, 760
Squash (see Cucurbits)
Statice (Limonium sinuatum): damping-off, caused by Botrytis cinerea, controlled with Clorox soak, benomyl soak or hot water treatment, 969
Stem and fruit disease: of papaya in Barbados, caused by Corynespora cassiicola, 404
Stemphylium botryosum: 592; 946
Stereum purpureum: 25; 192; 473
Stock (Matthiola incana): turnip mosaic virus found on annual stock or gilliflower in Argentina, 379
Stone fruits (see also Prunus): fungicides in hot water gave better control of Monilinia

fructicola on fruits than fungicidal dips in cold water, 62; control of M. fructicola on peach, plum and cherry fruit in New York with CW-524, a piperazine derivative fungicide, 457

Storage: of strawberry in CO_2-enriched atmospheres, 44; control of Botrytis storage rot of giant sequoia seedlings with fungicides, 67; cultivars of sugarbeet tested for resistance to storage rot, caused by Phoma betae, 204; reaction of onion cultivars to Botrytis brown stain, 210; of soybean affects seed quality, 770; of barley yellow dwarf virus by lyophilization, 862

---: of carrot: fungicide dips for controlling decay, 697; disease incidence in carrots influenced by amount of bacterial contamination in packinghouse waters and postharvest holding temperatures, 896

---: liquid nitrogen: storage of Macrophomina phaseolina sclerotia was an effective method over 9-month period, 478; longevity of rust spores in liquid nitrogen, 793

Strains: new strain of Xanthomonas phaseoli from Uganda, its virulence compared with strains from Colombia and Nebraska, 74; hypovirulent cultures of Endothia parasitica and their possible use for disease control in American chestnut, 225; gray and albino strains of Ceratocystis fagacearum and their recovery from inoculated trees, 254; severe strain of bean yellow mosaic virus is cause of chlorotic leaf spotting of yellow summer squash, 280; outbreak of strains of Cercospora beticola resistant to benzimidazole and thiophanate fungicides in northern Greece, 321

Strawberry: gray mold, caused by Botrytis cinerea, controlled with several fungicides, 31; quality and decay of California strawberries stored in CO_2-enriched atmospheres, 44; a new anthracnose caused by Colletotrichum dematium, found in Michigan, 445

Strophostyles helvola (quail pea): quail pea mosaic virus was isolated from quail pea and found to be a new member of the comovirus group, 311

Stylosanthes humilis: Colletotrichum leaf spot and stem canker on selections of this forage legume in Florida, 747

Sugarbeet: varietal resistance in sugarbeet cultivars to Phoma storage rot is more apparent at 20-25°C than at lower temperatures, 204; benomyl-resistant strains of Cercospora beticola caused severe losses to sugarbeets in Greece in 1972, 321; control of Heterodera schachtii on sugarbeet with aldicarb, 942

Sugarcane: hot air therapy at 54°C for 8 hr killed red rot fungus without damaging the buds, 220; leaf scald, inoculation of sugarcane plants with Xanthomonas albilineans quantitatively to screen for varietal resistance, 582; smut disease, its occurrence in Hawaii and efforts to control it, 966; ratoon stunting disease diagnosis by light microscopy, 987

Sugar content: nitrogen fertilization and mowing height affect sugar content in bluegrass leaves and susceptibility to leaf spot, 544

Sunflower (Helianthus annuus): 59 artificially inoculated sunflower selections were rated for susceptibility to the disease complex involving Fusarium moniliforme, Phymatotrichum omnivorum and Macrophomina phaseolina, 318; the confectionery, birdseed and oilseed varieties were evaluated for resistance to rust and Verticillium wilt, 524; basal gall, a new disease, caused by Plasmopara halstedii, 647

Survey: of production losses due to witches' broom of blueberry in the Maritime Provinces, 608

Survival: of Xanthomonas alfalfae in alfalfa debris and soil, 52; of sclerotia of Rhizoctonia solani, pathogenic to water hyacinth, 601; of Macrophomina phaseoli in corn and sorghum stalk residue as sclerotia, 873

Susceptibility: of tomato cultivars and lines to Rotylenchulus reniformis, 169; of tomato cultivars to leaf mold, caused by Cladosporium fulvum, under field conditions in Massachusetts, 179; of soybean to Meloidogyne naasi, found to be poor host, 207; of 'Marsh' grapefruit to greasy spot was influenced by the rootstock used, 294; to Diplodia maydis in stalks and ears of corn and the possible use of cob parenchyma cells for host-pathogen interaction studies, 308; of some apple cultivars to Physalospora obtusa infection, 676; multiplication and movement of Xanthomonas oryzae in susceptible and resistant hosts, 784; of peanut rotational crops tobacco, cotton and corn to Cylindrocladium crotalariae, 1035

Sweetpotato: breeding for root-knot resistance in sweetpotatoes is complicated by the variability in pathogenicity of Meloidogyne incognita races, 199; M. incognita were transmitted from infected bedding roots to the field via sprouts, but multiplication was not great enough to impair plant productivity of succeeding

crop, 371
Sycamore (Platanus occidentalis): ozone exposure caused significant reductions in growth of seedlings, 914
Symptomatology: of bean yellow mosaic virus, 280; of quail pea mosaic virus, 311; of bacterial wilt of cassava, 824
Symptom expression: of apple pucker disease with relation to temperature, 358
Synergism: of chloroneb and thiram in control of seed rot and damping-off in pea, 3

Take-all: of wheat, caused by Ophiobolus graminis, 187
Taphrina deformans: 361
Target spot: of tomato, in Haiti, 921
Taro (see Colocasia esculenta)
Temperature: hot air therapy, effective against red rot of sugarcane, 220; holding carrots in storage at 2°C for 2 or more days greatly reduced the incidence of decay from soft rot, 896
---: air: as a limiting factor in the development and spread of stem, leaf and stripe rusts of wheat in northern India, 8; temperature may be a limiting factor to disease development on pine of Sirococcus tip blight where winter and early spring months have monthly means below 9°C, 69; influence of temperature on recovery of Aspergillus flavus from soybean seed, 185; 20-25°C rather than 10-15° is better for selection for resistance in sugarbeet varieties to Phoma storage rot, 204; temperature, moisture and light play an important role in the sporulation of an isolate of race T of Helminthosporium maydis on a susceptible corn hybrid, 304; temperature effects on symptom expression of apple leaf pucker disease, 358; expression of petal symptoms on apple trees affected by leaf pucker, Jubilee ring-and-line pattern and Stayman blotch is apparently influenced by seasonal weather, 359; relationship of temperature and chemical concentration on phytotoxicity effect on cherry fruit, 428; high temperature coupled with high concentration of azinphosmethyl mixed with dodine or Difolatan caused greater toxicity to cherry fruit than any of chemicals alone, 428; relation of temperature and leaf wetness to development of leaf blight of wheat, 1000
---: hot water: the combination of fungicides with hot water for dips was more effective than unheated dip for control of brown rot of stone fruits, 62; hot water dips and heated DCNA controlled Pythium and Sclerotinia rots of snapbeans, 234; 15 minute dips at 118-119°F controlled Pratylenchus penetrans in dormant raspberry root cuttings, 492; hot water treatments of stored carrots reduced decay over untreated carrots, 697; hot water, for control of damping-off of statice, from Botrytis cinerea, 969
Texas: 81
Thuja orientalis (arborvitae): arborvitae blight controlled with benomyl sprays, 739
Tillage practices: not plowing under overwintered debris from infected corn fields is considered partially responsible for outbreak of Physoderma maydis in susceptible field corn lines in Illinois, 630
Tobacco: ozone injury occurred on tobacco indicator plants in South Dakota, possibly one of the earliest stages of plant-damaging air pollution ever recorded, 106; first report of lesion nematode on tobacco in Trinidad, 260; Corynespora leaf spot caused by C. cassiicola is reported for the first time on tobacco, in Nigeria, 375; black shank, report of first occurrence of race 1 of Phytophthora parasitica var. nicotianae in Georgia, 832; control of Sclerotinia in seedbeds with methyl bromide or DMTT fumigation, or drenches of benomyl or thiophanate methyl, 932; five flue-cured cultivars were highly susceptible to Cylindrocladium crotalariae, therefore, should probably not be used in peanut crop rotation, 1035
---: virus: three strains of potato virus Y found affecting flue-cured tobacco in the Southeastern United States, 200; veinbanding caused by potato virus Y, observed in flue-cured tobacco in 10 locations in North Carolina in 1972, 551; tobacco vein mottling virus has become widespread in Kentucky on burley tobacco, 841; all field-grown burley lines tested were susceptible to tobacco vein mottling virus, but showed range from tolerant to severe, 845
Tolerance: "horizontal resistance" in tomato against Fusarium wilt, 119; effect of Verticillium wilt on resistant, tolerant and susceptible tomato varieties, 122
Tomato: occurrence of Pythium disease with new seeding method in Florida with 1 : 1 mixture of vermiculite and peat moss together with seed and fertilizer, 260; branched broomrape (Orobanche ramosa) reduced yields of 'Manapal' tomato by 21-29% by direct measurement in California plots, 926
---: bacterial canker: caused by Corynebacterium michiganense found in California sprinkler-irrigated fields, 258;

Tomato (continued): new tomato variety H2990 showed high degree of resistance to C. michiganense, 584; C. michiganense was spread from diseased to healthy transplants by clipping with rotary mower, no control was obtained with tribasic copper sulfate, tetracycline HCl or streptomycin sulfate, 767; yield and mortality loss in H2990 (resistant) and H6 (susceptible) inoculated plants compared, 787; eradication from systemically infected tomato seed by a seed extraction process, with HCl, 974
---: bacterial speck, caused by Pseudomonas tomato, 258
---: bacterial spot, caused by Xanthomonas vesicatoria, found for first time on tomatoes in California, 258
---: early blight, segregation for horizontal resistance in tomato, 964
---: fruit rot: control of various tomato fruit rots with fungicides, 453; caused by Pythium aphanidermatum, 638; caused by two Pythium species in California, 1066
---: Fusarium wilt: symptoms reduced by three foliar sprays of acidified benomyl, 37; tolerance, or "horizontal resistance" of cultivar Homestead 24 showed little value in controlling the disease in Florida, 119; influenced by previous soil pasteurization with DD-MENCS or steam, 267; evaluation of tolerance and resistance to F. oxysporum lycopersici in 36 tomato varieties and the occurrence of a vertifolia effect, 724
---: late blight and leaf mold, control with fungicides caused damage to tomato fruits, 612
---: leafmold, cultivars show a range in susceptibility to leaf mold under field conditions in Massachusetts, 179
---: nematodes: reaction of root-knot nematode resistant cultivars with Mi resistant gene indicates that one or more genes other than Mi may affect root-knot nematode resistance in tomato, 141; genetic resistance to reniform nematode in tomato, 169; Meloidogyne incognita and Corynespora cassiicola were most serious pathogens observed in Haiti, 921; root-knot nematodes controlled with DD and DD-MENCS in Greece, 1033
---: Verticillium wilt, resistant, tolerant, and susceptible varieties were tested for their reaction to V. albo-atrum, 122
---: virus, tobacco mosaic virus was shown to decrease yield of fresh-market machine-harvest type tomatoes, 78
Toxin production: by Helminthosporium maydis as influenced by transmissible diseases, 420
Transmission: quail pea mosaic virus is transmitted by leaf beetles, 311; sprout transmission of Meloidogyne incognita to the field through infected sweetpotato bedding roots, 371; of the causal agent of isolates of Helminthosporium maydis and toxin specificity, 420; of maize dwarf mosaic virus, Schizaphis graminum more efficient than Aphis fabae or A. gossypii, 533; transmission and host range studies of broad bean mottle virus, 833
---: mechanical: effect of different buffers on the longevity at room temperature on relative infectivity of citrus exocortis virus from gynura, 598; of citrus exocortis virus, stem-cut inoculation with contaminated blade more efficient than needle puncture or leaf abrasion, 886
---: pollen transmission: of Prunus ringspot virus in prune and sour cherry orchards, 241; of pea seedborne mosaic virus in Pisum sativum, 248
---: seed: southern bean mosaic virus found to be seed-transmitted in two varieties of cowpea, 13; of tomato ringspot virus in raspberry, 431; of tomato ringspot virus in Gomphrena globosa, 433; of Pseudomonas in Pelargonium hortorum, 909
Trap-crop: disulfoton-treated pearl millet planted around cowpeas apparently decreased the number of cowpea mosaic-infected cowpeas within screened area compared with the unscreened control plants, 684
Trichoderma spp.: as a possible biological control to reduce the incidence of gray snow mold on turfgrass, 924
--- viride: as a biological control agent, applied by pruning shears controlled Stereum purpureum infection of pruning wounds on plum trees, 25
Tsuga heterophylla (western hemlock): contrasting effects of two herbicides on invasion by Fomes annosus in tree-injector wounds on young trees, 94
Turkey: 149
Turnip (Brassica rapa, see Crucifers
Tylenchorhynchus spp.: 891
--- dubius: 438
Typhula incarnata: 924

Ulmus americana (American elm, see Elm)
Ulmus parvifolia (Chinese elm, see Elm)
Urd bean (Phaseolus mungo): viral diseases that cause losses in India, 463; white fly-transmitted yellow mosaic of urd bean from West Pakistan, 800
Uredinopsis sp.: 765
Uromyces phaseoli: 793
Ustilago scitaminea: 966

Variability: of Meloidogyne incognita races in their pathogenicity to sweetpotatoes, 199
Vectors associated with: bean yellow mosaic virus, Macrosiphum euphorbiae, 280; bunchy top disease of papaya, Empoasca stevensi, 412; mung bean yellow mosaic virus, Bemisia tabaci, 463; Dutch elm disease, Scolytus multistriatus, 511; 701; maize dwarf mosaic virus, by Aphis fabae, Aphis gossypii, and Schizaphis graminum, 533; pea seedborne mosaic virus, in Washington, seven aphid species, Acyrthosiphon pisum, Dactynotus escalantii, Macrosiphum euphorbiae, M. rosae, Myzus persicae, Ovatus crataegarius, and Rhopalosiphum padi were efficient vectors, 552; lettuce mosaic virus, Acyrthosiphon pisum, 688; tobacco vein mottling virus, Myzus persicae and Macrosiphum euphorbiae, 841
Veitchia merrillii (Christmas palm): susceptible to lethal yellowing disease in Florida, 861
Ventilation: lack of ventilation and excess moisture appeared to increase disease incidence of Santolina blight of gray and green Santolinas, 1072
Venturia inaequalis: 560; 817
Vermont: 338; 394
Verticillium albo-atrum: 122; 284; 403; 460; 524; 928
--- dahliae: 28; 735; 928
Vertifolia effect: to 36 tomato varieties tested for resistance and tolerance to Fusarium oxysporum lycopersici, 724
Vicia villosa (hairy vetch): pea seedborne mosaic virus will overwinter effectively in seedlings of hairy vetch under experimental field conditions, 349
Vigna sinensis (see Cowpea)
Virginia: 182
Virgin soil: a new seeding method of addition of vermiculite and peat moss with tomato seed and fertilizer to virgin sandy soils in Florida resulted in a stunting and root-rotting, caused by Pythium spp., 260
Virulence: comparative virulence of Xanthomonas phaseoli strains from Uganda, Colombia and Nebraska, 74; the age of Septoria nodorum inoculum greatly affects its ability to infect wheat, 363; variation in colony morphology and its relation to virulence in Xanthomonas oryzae, 955
Virus: of several grasses, including corn and sorghum, reported from Israel for the first time, 125; virus found in black locust in Arkansas resembles tomato spotted wilt virus, 153; viruslike symptoms in ash found to vary from year to year on the same trees over a 6-year period of study, 396; virus diseases of some warm weather pulse crops in India, 463; unidentified virus produced bacilliform particles in the cortex cells of fruits of Cucumis melo, 649; of peppers in Florida, 991; of blackjack oak and black oak, viruslike flexuous rods found in cytoplasm of leaves of trees with ringspot symptoms, 1029
---: barley stripe mosaic virus, aphids, a leafhopper and a grasshopper failed to transmit the virus, 639
---: barley yellow dwarf, preservation by lyophilization, 862
---: bean pod mottle virus of soybeans: 811; effect on yield components and morphology of soybeans in relation to soil water regimes, 1050
---: bean yellow mosaic virus, found to be cause of chlorotic leaf spotting of yellow summer squash, 280
---: broad bean mottle virus, transmission and host range studies, 833
---: bromegrass mosaic, identified from South Dakota in corn and sorghum, 373
---: citrus exocortis virus: extraction and mechanical transmission of citrus exocortis virus in gynura, 598; mechanical transmission, 886; monoembryonic seedlings of Arizona 861 selection of 'Etrog' citron show uniform response to the virus, 1010
---: chlorotic leaf spot, distribution in apple budsticks, 865
---: corn stunt, spiral-shaped microorganisms occur naturally in corn infected with corn stunt virus in Mexico, 333
---: cowpea mosaic virus, cowpeas planted within a pearl millet trap-crop (treated with disulfoton insecticide) developed fewer virus-infected cowpea plants than control plants, 684
---: cucumber mosaic virus, found in Nandina domestica in South Carolina, 917
---: cymbidium mosaic virus, control of powdery mildew on Cassia occidentalis with benomyl did not affect the plant's local lesion response to the virus, 475
---: Hydrangea ringspot virus, the cause of phyllody of florists' Hydrangea, 300
---: leaf pucker: in McIntosh apple as affected by temperature, 358; leaf pucker, Jubilee ring-and-line pattern, and Stayman blotch, blossom symptoms, 359
---: lettuce mosaic virus, natural occurrence in Pisum sativum in New York, 688
---: maize dwarf mosaic: transmission of virus by Aphis fabae, A. gossypii, and Schizaphis graminum compared, S. graminum is more efficient than either of others, 533; found in South Dakota in

Virus (continued): 1972, not previously reported from that State, 373
---: maize streak virus, an additional note on viruslike particles associated with diseased material from Africa, 414
---: mung bean yellow mosaic virus, its symptomatology, transmission, host range, incidence, losses and control in India, 463
---: pea seedborne mosaic virus: further studies on seed transmission, 248; its ability to overwinter in hairy vetch is a danger if hairy vetch growing in close proximity to pea-growing areas should become infected with the virus, 349; potential aphid vectors in Washington State, 552
---: pear vein yellows virus, incidence, systemic nature, and spread in the Yakima Valley, Washington, 483
---: potato virus Y (PVY), three distinct strains can be differentiated on root-knot nematode susceptible and resistant flue-cured tobacco varieties in the Southeastern United States, 200; a new outbreak on tobacco in North Carolina, 551; potato virus Y on pepper, 991
---: Prunus ringspot virus, pollen transmission in prune and sour cherry orchards, 241
---: quail pea mosaic virus, differs in host range and symptomatology from bean pod mottle and cowpea mosaic viruses, but is related serologically to other members of the Comovirus group, 311
---: Satsuma dwarf virus, found for the first time in Satsuma mandarin in the Aegean Coast of Turkey, 149
---: southern bean mosaic, effect of plant variety and infection age on the presence of the virus in floral parts and unripe seeds of Vigna sinensis, 13
---: tobacco etch virus of pepper, 991
---: tobacco mosaic, caused decrease in yield of fresh-market machine-harvest tomatoes, with earlier infection causing greater reduction in yield than later infections, 78; tobacco mosaic virus on pepper, 991
---: tobacco ringspot virus: was recovered from 7% of symptomatic transplanted ash trees showing viruslike symptoms, 396; infection in soybeans detected by aerial photography, 471; effects of infection in soybean fields in Louisiana were about twice as severe as those for bean pod mottle virus, 811
---: tobacco vein mottling virus, on burley tobacco in Kentucky, 841; varietal reaction of field-grown burley cultivars, 845
---: tomato ringspot virus: seed-transmitted in raspberry, 431; is seed transmitted in Gomphrena globosa, 433
---: turnip mosaic, found on annual stock (Matthiola incana) in Argentina, 379; turnip mosaic virus and cauliflower mosaic viruses identified in turnip and mustard plants in Georgia, 978
---: urd bean mosaic mottle, urd bean leaf crinkle, and urd bean leaf curl, in India, 463
---: wheat streak mosaic: attempt was made to determine which chromosome in rye carries the gene(s) for reaction to wheat streak mosaic virus, 6; recovered from corn and sorghum in South Dakota, 373
Viscum album: 904

Walnut (Juglans nigra): seedlings display ozone tolerance, 914
Washington: 164; 167; 483; 522; 552; 865
Water: soil water, as stress factor coupled with bean pod mottle virus infection reduced soybean yields more than infection alone, 1050
Water hyacinth (Eichhornia crassipes): sclerotia of an isolate of Rhizoctonia solani pathogenic to water hyacinth survived for 26 months in lake water, 601
Watery soft rot: of bean, caused by Sclerotinia sclerotiorum, controlled with post-harvest dip in hot water or DCNA, 234
Wax treatment: of mangos, following hot dip treatment with benomyl controlled shriveling of fruits if applied immediately after treatment, 173
Weather: root rot and death of ohia trees in Hawaii from Phytophthora cinnamomi was related to high rainfall and poor drainage, but not to ohia decline, 383; warm, wet spring weather and conservation tillage possibly responsible for outbreak of Physoderma maydis infection in field corn in Illinois, 630
Weeds: as potential reservoirs of disease, Hordeum leporinum is host of Septoria halophila in Australia, 367
West Indies: 260
West Pakistan: 505; 800
West Virginia: 1055
Wheat: reaction of rye addition lines in wheat to wheat streak mosaic virus, 6; Helminthosporium blight incidence in India, and results of screening varieties for resistance, 278; proved to be a good host of Pratylenchus penetrans, 728; rate of increase of leaf blight caused by Alternaria triticina in relation to temperature, leaf wetness and host age, 1000
---: rust: epidemiology of stem, leaf and stripe rusts in northern India, 8; leaf rust and powdery mildew were the most prevalent diseases of wheat in Michigan

over a 4-year period, eight others were also common, 187; stem rust epidemic potential in 1972, 434; virulence of wheat leaf rust in North Dakota, in 1972, 449; leaf and stem rust control by seed treatment with oxycarboxin is apparent at initial stages of rust development, but late disease development obscured effects of early control on yield of wheat, 567; BT applied through irrigation water or as spray controlled leaf rust, in India, 572; early application of BT fungicide effected best control of leaf rust, 653; prevalence of races of stem rust in the U. S. in 1972, 880

---: Septoria: relative prevalence of Septoria species on spring wheat in Minnesota, 99; age of cultures of S. nodorum for inoculum preparation is important, 4- to 6-day-old cultures more effective than older cultures, 363

Whetzelinia sclerotiorum (= Sclerotinia sclerotiorum): 164; 234; 770; 932; 1070

White mold: of Great Northern beans, wider row spacing and determinate growth reduced severity, 1070

Wilt: bacterial: of carnation, caused by Erwinia chrysanthemi, in New Zealand, 136; Goss's bacterial wilt, also called leaf freckles of corn, caused by undescribed Corynebacterium species, 291; of cassava, caused by Xanthomonas manihotis, in Nigeria, 824

---: Fusarium: of cotton, yield reduction has remained fairly constant over a 7-year period, 325; of variegated Pyracantha, controlled with benomyl, 389

---:---: of tomato: three foliar sprays of acidified benomyl reduced Fusarium wilt symptoms in inoculated tomato plants by about 50%, 37; failure of horizontal resistance in Florida test, 119; more severe wilt developed in inoculated plants that were planted in pasteurized soil than in unpasteurized soil, 267; variety resistance and tolerance in 36 cultivars, 724

---: oak wilt: caused by Ceratocystis fagacearum, a study to determine the compatibility and color types that can be recovered from mixed inoculum used to inoculate scarlet oak trees, 254; the imperfect stage of C. fagacearum was isolated from water oak near La Grange, North Carolina approx. 200 miles east of closest known center in N. C., and 170 miles northeast of infection in S. C., 325

---: Verticillium wilt: of tomato, effect on resistant, tolerant, and susceptible tomato varieties, 122; of cotton, on national basis, losses have decreased in a 7-year period, 325; of cotton, the high concentration of CW-524 necessary to effect control of Verticillium wilt would appear to eliminate it as an effective systemic fungicide for this disease, 460; resistance in sunflower varieties, to wilt and rust, 524; in Rhode Island-grown potatoes developed in North Dakota, 928

Wind: aerial detection of tobacco ringspot virus in soybean fields suggests the probable involvement of windborne insect vectors in virus spread, 471

Wisconsin: 349; 473

Witches' broom: of blueberry, survey of production losses from, 608

Wyoming: 576

Xanthomonas: 284

--- albilineans: 582; 1040

--- alfalfae: 52

--- malvacearum: 403

--- manihotis: 824

--- nigromaculans f. sp. zinniae: 555; 1020

--- oryzae: 227; 537; 784; 955

--- phaseoli: 74; 111

--- vesicatoria: 258

Xiphinema americanum: 416

--- index: 115

Yellow birch (see Betula alleghaniensis)

Yellow poplar (Liriodendron tulipifera): seedlings exposed to ozone showed little growth change but leaf drop occurred, 914

Yield: of soybean, effect of Cephalosporium gregatum on soybeans in Illinois, 853

Yield increase: of alfalfa hay in Minnesota with zinc + maneb, 353

Yield loss: to tomatoes (fresh-market machine-harvest) from infection by tobacco mosaic virus, 78; of cotton, loss from nematodes and boll rots from 1965-1971 has increased on a national basis, loss from Verticillium wilt has decreased, 325; of cotton by parasitic diseases in 1972, 403; of blueberry, due to witches' broom, in the Maritime Provinces, 608; effects of tobacco ringspot virus infection in soybeans in Louisiana were almost twice as severe as those from bean pod mottle virus as measured by reduction in yields, 811; in tomatoes, from Orobanche ramosa, 926

Zinnia: bacterial leaf spot, caused by Xanthomonas nigromaculans f. sp. zinniae, found on cultivars of zinnia in Columbus, Ohio in 1972, first report in the United States, 555; seed dissemination of Xanthomonas nigromaculans f. sp. zinniae in infected zinnia seed indicated, 1020

Prepared by Marguerite G. Benedict